

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/82, 15/29, C07K 14/415, C12N 5/10, A01H 1/00	A1	(11) International Publication Number: WO 00/32799 (43) International Publication Date: 8 June 2000 (08.06.00)
(21) International Application Number: PCT/US99/28103 (22) International Filing Date: 24 November 1999 (24.11.99) (30) Priority Data: 60/109,892 25 November 1998 (25.11.98) US (71) Applicant: CALGENE LLC [US/US]; 1920 Fifth Street, Davis, CA 95616 (US). (72) Inventor: CHAUDHURI, Sumita; 1125 Juniper Place, Davis, CA 95616 (US). (74) Agent: SCHWEDLER, Carl, J.; Patent Department Central, Monsanto/G.D. Searle, P.O. Box 5110, Chicago, IL 60680-5110 (US).		(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHODS FOR TRANSFORMING PLASTIDS (57) Abstract An improved method is provided for the transformation of a plant cell plastid. The improved method allows for the increased efficiency of the foreign DNA penetrating the plastid membrane. The method generally involves the use of a plant tissue source having an altered plastid morphology in plastid transformation methods. The present invention finds use in plastid transformation methods for a wide variety of plant species.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHODS FOR TRANSFORMING PLASTIDS

5

FIELD OF THE INVENTION

This invention relates to the application of genetic engineering techniques to plants. More specifically, the invention relates to methods for the transformation of
10 plant cell plastids.

BACKGROUND

15 The plastids of higher plants are an attractive target for genetic engineering. Plant plastids (chloroplasts, amyloplasts, elaioplasts, chromoplasts, etc.) are the major biosynthetic centers that in addition to photosynthesis are responsible for production of industrially important compounds such as amino acids, complex carbohydrates, fatty acids, and pigments. Plastids are derived from a common precursor known as a
20 proplastid and thus the plastids present in a given plant species all have the same genetic content. Plant cells contain 500-10,000 copies of a small 120-160 kilobase circular genome, each molecule of which has a large (approximately 25kb) inverted repeat. Thus, it is possible to engineer plant cells to contain up to 20,000 copies of a particular gene of interest which potentially can result in very high levels of foreign
25 gene expression.

Current plastid transformation methods are inefficient, as such there is need for constructs and methods which improve plastid transformation.

SUMMARY OF THE INVENTION

30

By this invention, methods which allow for the improved transformation of a foreign DNA into plant cell plastids are provided. Such methods generally involve

utilizing a plant tissue source which contains cells with an altered plastid morphology in the transformation methods. The alteration in the plant plastid morphology includes, *inter alia*, plastid size and number. By utilizing tissue derived from such plants in plastid transformation methods, efficiency of transformation of a foreign
5 DNA into the plant cell plastid may be increased.

As exemplified herein, constructs useful for genetic engineering of plant cells to provide for a method of increasing plastid transformation efficiency are provided. The constructs include nucleic acid sequences coding for protein sequences involved in controlling division of plant cell organelles. The expression of such nucleic acid
10 sequences in a plant cell provides for an altered number and/or size of the chloroplasts within the host cell.

DNA sequences, also referred to herein as polynucleotides, for use in transformation contain an expression construct comprising a promoter region which is functional in a plastid, and a DNA sequence encoding a gene involved in controlling
15 the division of plant cell organelles.

Methods for the use of transformed plants with altered plastid morphology are described. Such methods include plant breeding or transformation methods to provide plant cells having both the nuclear and plastid constructs.

The present invention also provides methods for increasing the efficiency of
20 chloroplast transformation. The method generally comprises transforming the plastids of a plant tissue which has been modified to have an altered number and/or size of plastids contained within the plant cell.

The present invention also provides a mechanism for enhancing the efficiency of chloroplast transformation in plant species.

25 The present invention also provides methods for improving the selectability of plant comprising, transforming a plant cell source having an altered plastid morphology with a construct comprising a promoter functional in a plant cell plastid operably associated with a nucleic acid sequence encoding a selectable marker. Selectable markers of interest in the present invention include herbicide tolerance
30 genes such as glyphosate tolerance genes, and antibiotic resistance genes. Glyphosate tolerance genes include the CP4 gene from *Agrobacterium*.

Another aspect of the present invention provides methods for preparing a plant cell source with increased plastid transformation efficiency comprising, transforming a plant cell with a construct comprising a promoter functional in plant cell operably associated with a nucleic acid sequence encoding a FtsZ protein.

5 Also considered part of this invention are the plants and plant cells obtained using the methods described herein.

DESCRIPTION OF THE FIGURES

10 Figure 1 provides an amino acid sequence alignment of the Arabidopsis FtsZ1 (SEQ ID NO:2), the Brassica FtsZ1 (SEQ ID NO:6), the tobacco FtsZ1 (SEQ ID NO:9), the Soybean FtsZ1 (SEQ ID NO:72) and the corn FtsZ1 (SEQ ID NO:73) protein sequences.

15

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the subject invention, methods are provided which allow for the improved transformation of a foreign DNA into plant cell plastids. Such methods generally involve utilizing a plant cell source which contains an altered plant plastid morphology. By utilizing tissue derived from such plants in plastid transformation methods, efficiency of transformation of a foreign DNA into the plant cell plastid can be increased.

25 In one embodiment of the instant invention, plant tissue containing altered plant plastid morphology is used for plastid transformation methods. Such alterations in plant plastid morphology include, but are not limited to, alterations in the plastid size, shape and number in respect to a wild-type plastid morphology from the target plant cell. In general, a wild-type plastid morphology consists of small, round organelles contained within the plant cell, depending on the species. Furthermore, a plant cell typically contains between about 50 and about 100 plastids.

30 The plant tissue source used in plastid transformation methods of the present invention contains an increase in the size of the plastids contained in the plant cells.

Such increases in the size of the plastids provides for a larger surface area for the foreign DNA to penetrate the plastid membrane during transformation.

The large plastids preferably contain approximately the same number of plastid genomes as would be contained in corresponding number of wild-type plastids. For example, in a wild-type plant cell containing 100 plastids per cell and 100 copies of the plastid genome in each plastid (a total of 10,000 copies of the plastid genome per cell), the corresponding mutant tissue source would preferably contain about the same number of plastid genomes, only contained in one, or several large plastid(s).

Alternatively, a plant tissue source with an increased number of plastids, with a corresponding reduced size, can also find use in the plastid transformation methods of the present invention.

As is understood in the art, additional methods for obtaining plants with alterations in the plastid size and number are known. The skilled artisan will recognize that a number of methods are available for providing for an alteration in plastid cell division. Such methods are described, for example, by Strepp, *et al.* (1998) *Proc. Natl. Acad. Sci. USA*, 95:4368-4373.

Cell division, also referred to as cytokinesis, has been the focus of studies in many organisms such as bacterial, fungal, and animal cells. Division of bacterial cells occurs through the formation of an FtsZ ring (also referred to as a Z ring) at the site of division (Lutkenhaus, *et al.* (1997) *Ann. Rev. Biochem.* 66:93-116). The positioning and formation of the Z ring acts to further drive septation (cytokinesis). The ring is composed of a tubulin-like FtsZ protein which has GTPase activity. Mutations in the *ftsZ* gene in *E. coli* leads to the production of a temperature-sensitive filaments with regularly spaced nucleoids at certain temperatures (Lutkenhaus (1992) In *Prokaryotic Structure and Function: A New Perspective*, ed. S Mohan, C Dow, pp 123-152. Cambridge: Cambridge Univ. Press). Such mutations in bacteria leads to the inability to divide correctly.

The plant cell plastid as well as the mitochondria are derived from prokaryotic ancestors, and thus, the division apparatus of these organelles resembles that of bacteria. Recently, identification of *ftsZ* related sequences in *Arabidopsis* and *Physcomitrella patens* have been reported (Osteryoung, *et al.* (1995) *Nature*, 376:473-

474; and Strepp, *et al.* (1998), *supra*). The protein encoded by the *Arabidopsis ftsZ* gene was found to be imported into the chloroplast and was therefore speculated to be a component of the plastid division machinery (Osteryoung, *et al.* (1995), *supra*). More recently, the involvement of FtsZ in plastid division was directly demonstrated.

5 The disruption of the *ftsZ* gene in a lower plant, *Physcomitrella patens*, impeded plastid division, thereby giving rise to mutant cell lines with one or a few large plastids (Strepp, *et al.* (1998), *supra*).

The use of plants with an altered number and/or size of plastids containing one or few large plastids could therefore be used as targets for plastid transformation of any plant species. Such plants containing an altered size and/or number can be

10 obtained using various methods, including mutagenesis, antisense suppression, or co-suppression. Methods for the mutagenesis of plant genomes are well known in the art, and include chemical, such as ethylmethane sulfonate (EMS) and nitrosoguanidine (NTG), as well as physical mutagenesis methods such as fast neutron bombardment.

15 Other means for obtaining a plant source with an alteration in the size and/or number of plastids contained in the cell are also contemplated. For example, tissue for use in the transformation methods of the present invention can be obtained from plants grown in culture conditions which provide for such altered plastid content. For example, tissue obtained from plants grown *in vitro* under culture conditions in which

20 inhibitors of bacterial cell division, such as 5,5'-bis-(8-anilino-1-naphtalenesulfonate) (Yu, *et al.* (1998) *J. Biol Chem.* 273:10216-10222), are present, can be utilized as a cell source for the plastid transformation methods of the present invention.

In a preferred embodiment, such plants containing cells with an alteration in the size and/or number of plastids are generated by anti-sense expression of the FtsZ

25 gene. Once plastid transformation is achieved and homoplasmic plants are identified, the anti-sense transgene can be eliminated by out-crossing and the wild-type condition of 50 to 100 plastids per cell restored. Similarly, plants regenerated from plastid transformed tissue containing an altered number and/or size of plastids from mutations can also be reverted to the wild-type plastid conditions using such out-

30 crossing methods.

In the case of the use of culture conditions for obtaining plant cells with an altered number and/or size of plastids, wild-type plastids can be obtained by releasing the tissue from such culture conditions.

In another embodiment of the present invention, novel nucleic acid sequences
5 are provided which encode proteins related to proteins involved in bacterial cell and plastid division.

In particular, novel nucleic acid sequences from *Arabidopsis*, soybean, corn, Brassica are provided which encode FtsZ related proteins. Such nucleic acid sequences find use in the preparation of DNA constructs. Such constructs find use in
10 the production of plants with an altered number and/or size of chloroplasts.

The skilled artisan will recognize that other DNA sequences useful for the production of plants with an altered number and/or size of chloroplasts are available in the art. The sequences include but are not limited to, *ftsA*, *ftsL*, *ftsI*, *ftsQ*, *ftsN*, *ftsW*, *ftsK* (Lutkenhaus, *et al.* (1997) *supra*), and the *arc* genes (Pyke, *et al.* (1992) *Plant Physiol.* 99:1005-1008; Pyke *et al.* (1994) *Plant Physiol.* 104:201-207; and
15 Pyke (1997) *Am. J. Botany* 84:1017-1027).

In order to obtain additional *ftsZ* sequences, a genomic or other appropriate library prepared from the candidate plant source of interest can be probed with conserved sequences from one or more plant and/or bacterial *ftsZ* sequence(s) to
20 identify homologically related sequences. Positive clones can be analyzed by restriction enzyme digestion and/or sequencing. When a genomic library is used, one or more sequences can be identified providing both the coding region, as well as the transcriptional regulatory elements of the *ftsZ* gene from such plant source. Probes can also be considerably shorter than the entire sequence. Oligonucleotides can be
25 used, for example, but should be at least about 10, preferably at least about 15, more preferably at least 20 nucleotides in length. When shorter length regions are used for comparison, a higher degree of sequence identity is required than for longer sequences. Shorter probes are often particularly useful for polymerase chain reactions (PCR), especially when highly conserved sequences can be identified. (See, Gould, *et*
30 *al.*, *PNAS USA* (1989) 86:1934-1938.)


When longer nucleic acid fragments (>100 bp) are employed as probes, especially when using complete or large cDNA sequences, one can still screen with

moderately high stringencies (for example using 50% formamide at 37°C with minimal washing) in order to obtain signal from the target sample with 20-50% deviation, i.e., homologous sequences. (For additional information regarding screening techniques see Beltz, *et al.*, *Meth. Enzymology* (1983) 100:266-285).

5 Another aspect of the present invention relates to isolated FtsZ polynucleotides. The polynucleotide sequences of the present invention include isolated polynucleotides that encode the polypeptides of the invention having a deduced amino acid sequence selected from the group of sequences set forth in the Sequence Listing and to other polynucleotide sequences closely related to such
10 sequences and variants thereof.

The invention provides a polynucleotide sequence identical over its entire length to each coding sequence as set forth in the Sequence Listing. The invention also provides the coding sequence for the mature polypeptide or a fragment thereof, as well as the coding sequence for the mature polypeptide or a fragment thereof in a
15 reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, pro-, or prepro- protein sequence. The polynucleotide can also include non-coding sequences, including for example, but not limited to, non-coding 5' and 3' sequences, such as the transcribed, untranslated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns,
20 polyadenylation signals, and additional coding sequence that encodes additional amino acids. For example, a marker sequence can be included to facilitate the purification of the fused polypeptide. Polynucleotides of the present invention also include polynucleotides comprising a structural gene and the naturally associated sequences that control gene expression.

25 The invention also includes polynucleotides of the formula:



wherein, at the 5' end, X is hydrogen, and at the 3' end, Y is hydrogen or a metal, R₁ and R₃ are any nucleic acid residue, n is an integer between 1 and 3000, preferably between 1 and 1000 and R₂ is a nucleic acid sequence of the invention, particularly a
30 nucleic acid sequence selected from the group set forth in the Sequence Listing and preferably SEQ ID NOs: 1, 3, 5, 7, 8, and 10-31. In the formula, R₂ is oriented so that its 5' end residue is at the left, bound to R₁, and its 3' end residue is at the right, bound to

R₃. Any stretch of nucleic acid residues denoted by either R group, where R is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

The invention also relates to variants of the polynucleotides described herein that encode for variants of the polypeptides of the invention. Variants that are
5 fragments of the polynucleotides of the invention can be used to synthesize full-length polynucleotides of the invention. Preferred embodiments are polynucleotides encoding polypeptide variants wherein 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues of a polypeptide sequence of the invention are substituted, added or deleted, in any combination. Particularly preferred are substitutions, additions, and deletions
10 that are silent such that they do not alter the properties or activities of the polynucleotide or polypeptide.

Further preferred embodiments of the invention that are at least 50%, 60%, or 70% identical over their entire length to a polynucleotide encoding a polypeptide of the invention, and polynucleotides that are complementary to such polynucleotides.
15 More preferable are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding a polypeptide of the invention and polynucleotides that are complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length are particularly preferred, those at least 95% identical are especially preferred. Further, those with at
20 least 97% identity are highly preferred and those with at least 98% and 99% identity are particularly highly preferred, with those at least 99% being the most highly preferred.

Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptides encoded by the polynucleotides set forth in the Sequence Listing.
25

The invention further relates to polynucleotides that hybridize to the above-described sequences. In particular, the invention relates to polynucleotides that hybridize under stringent conditions to the above-described polynucleotides. As used herein, the terms "stringent conditions" and "stringent hybridization conditions" mean
30 that hybridization will generally occur if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising 50% formamide,

5x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/milliliter denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at approximately 65°C. Other hybridization and wash conditions are well known and are exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY (1989), particularly Chapter 11.

The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in the Sequence Listing under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers as described herein.

As discussed herein regarding polynucleotide assays of the invention, for example, polynucleotides of the invention can be used as a hybridization probe for RNA, cDNA, or genomic DNA to isolate full length cDNAs or genomic clones encoding a polypeptide and to isolate cDNA or genomic clones of other genes that have a high sequence similarity to a polynucleotide set forth in the Sequence Listing. Such probes will generally comprise at least 15 bases. Preferably such probes will have at least 30 bases and can have at least 50 bases. Particularly preferred probes will have between 30 bases and 50 bases, inclusive.

The coding region of each gene that comprises or is comprised by a polynucleotide sequence set forth in the Sequence Listing may be isolated by screening using a DNA sequence provided in the Sequence Listing to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to identify members of the library which hybridize to the probe. For example, synthetic oligonucleotides are prepared which correspond to the FtsZ EST sequences. The oligonucleotides are used as primers in polymerase chain reaction (PCR) techniques to obtain 5' and 3' terminal sequence of FtsZ genes. Alternatively, where oligonucleotides of low degeneracy can be prepared from particular FtsZ

peptides, such probes may be used directly to screen gene libraries for FtsZ gene sequences. In particular, screening of cDNA libraries in phage vectors is useful in such methods due to lower levels of background hybridization.

Typically, a FtsZ sequence obtainable from the use of nucleic acid probes will
5 show 60-70% sequence identity between the target FtsZ sequence and the encoding sequence used as a probe. However, lengthy sequences with as little as 50-60% sequence identity may also be obtained. The nucleic acid probes may be a lengthy fragment of the nucleic acid sequence, or may also be a shorter, oligonucleotide probe. When longer nucleic acid fragments are employed as probes (greater than about 100
10 bp), one may screen at lower stringencies in order to obtain sequences from the target sample which have 20-50% deviation (i.e., 50-80% sequence homology) from the sequences used as probe. Oligonucleotide probes can be considerably shorter than the entire nucleic acid sequence encoding an FtsZ enzyme, but should be at least about 10, preferably at least about 15, and more preferably at least about 20 nucleotides. A
15 higher degree of sequence identity is desired when shorter regions are used as opposed to longer regions. It may thus be desirable to identify regions of highly conserved amino acid sequence to design oligonucleotide probes for detecting and recovering other related FtsZ genes. Shorter probes are often particularly useful for polymerase chain reactions (PCR), especially when highly conserved sequences can be identified.
20 (See, Gould, *et al.*, *PNAS USA* (1989) 86:1934-1938.).

Another aspect of the present invention relates to FtsZ polypeptides. Such polypeptides include isolated polypeptides set forth in the Sequence Listing, as well as polypeptides and fragments thereof, particularly those polypeptides which exhibit FtsZ activity and also those polypeptides which have at least 50%, 60% or 70%
25 identity, preferably at least 80% identity, more preferably at least 90% identity, and most preferably at least 95% identity to a polypeptide sequence selected from the group of sequences set forth in the Sequence Listing, and also include portions of such polypeptides, wherein such portion of the polypeptide preferably includes at least 30 amino acids and more preferably includes at least 50 amino acids.

30 "Identity", as is well understood in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence

relatedness between polypeptide or polynucleotide sequences, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods including, but not limited to, those described in *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York (1988);

5 *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M. and Griffin, H.G., eds., Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press (1987); *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., Stockton Press, New York (1991); and

10 Carillo, H., and Lipman, D., *SIAM J Applied Math*, 48:1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available programs. Computer programs which can be used to determine identity between two sequences include, but are not limited to, GCG (Devereux, J., et al., *Nucleic Acids*

15 *Research* 12(1):387 (1984); suite of five BLAST programs, three designed for nucleotide sequences queries (BLASTN, BLASTX, and TBLASTX) and two designed for protein sequence queries (BLASTP and TBLASTN) (Coulson, *Trends in Biotechnology*, 12: 76-80 (1994); Birren, et al., *Genome Analysis, I*: 543-559 (1997)). The BLAST X program is publicly available from NCBI and other sources (*BLAST*

20 *Manual*, Altschul, S., et al., NCBI NLM NIH, Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.*, 215:403-410 (1990)). The well known Smith Waterman algorithm can also be used to determine identity.

Parameters for polypeptide sequence comparison typically include the following:

- 25 Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)
 Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, *Proc. Natl. Acad. Sci USA* 89:10915-10919 (1992)
 Gap Penalty: 12
 Gap Length Penalty: 4
- 30 A program which can be used with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison Wisconsin. The above

parameters along with no penalty for end gap are the default parameters for peptide comparisons.

Parameters for polynucleotide sequence comparison include the following:

Algorithm: Needleman and Wunsch, J. Mol. Biol. 48:443-453 (1970)


5 Comparison matrix: matches = +10; mismatches = 0

Gap Penalty: 50

Gap Length Penalty: 3

A program which can be used with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison Wisconsin. The above
10 parameters are the default parameters for nucleic acid comparisons.

The invention also includes polypeptides of the formula:



wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal, R_1 and R_3 are any amino acid residue, n is an integer between 1
15 and 1000, and R_2 is an amino acid sequence of the invention, particularly an amino acid sequence selected from the group set forth in the Sequence Listing and preferably SEQ ID NOs: 2,4,6, and 9. In the formula, R_2 is oriented so that its amino terminal residue is at the left, bound to R_1 , and its carboxy terminal residue is at the right, bound to R_3 . Any stretch of amino acid residues denoted by either R group, where R
20 is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

Polypeptides of the present invention include isolated polypeptides encoded by a polynucleotide comprising a sequence selected from the group of a sequence contained in the Sequence Listing set forth herein .

25 The polypeptides of the present invention can be mature protein or can be part of a fusion protein.

Fragments and variants of the polypeptides are also considered to be a part of the invention. A fragment is a variant polypeptide which has an amino acid sequence that is entirely the same as part but not all of the amino acid sequence of the
30 previously described polypeptides. The fragments can be "free-standing" or comprised within a larger polypeptide of which the fragment forms a part or a region, most preferably as a single continuous region. Preferred fragments are biologically

active fragments which are those fragments that mediate activities of the polypeptides of the invention, including those with similar activity or improved activity or with a decreased activity. Also included are those fragments that antigenic or immunogenic in an animal, particularly a human.

5 Variants of the polypeptide also include polypeptides that vary from the sequences set forth in the Sequence Listing by conservative amino acid substitutions, substitution of a residue by another with like characteristics. In general, such substitutions are among Ala, Val, Leu and Ile; between Ser and Thr; between Asp and Glu; between Asn and Gln; between Lys and Arg; or between Phe and Tyr.

10 Particularly preferred are variants in which 5 to 10; 1 to 5; 1 to 3 or one amino acid(s) are substituted, deleted, or added, in any combination.

Variants that are fragments of the polypeptides of the invention can be used to produce the corresponding full length polypeptide by peptide synthesis. Therefore, these variants can be used as intermediates for producing the full-length polypeptides
15 of the invention.

The polynucleotides and polypeptides of the invention can be used, for example, in the transformation of host cells, such as plant host cells, as further discussed herein.

The invention also provides polynucleotides that encode a polypeptide that is a
20 mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids within the mature polypeptide (for example, when the mature form of the protein has more than one polypeptide chain). Such sequences can, for example, play a role in the processing of a protein from a precursor to a mature form, allow protein transport, shorten or lengthen protein half-life, or facilitate manipulation of the protein
25 in assays or production. It is contemplated that cellular enzymes can be used to remove any additional amino acids from the mature protein.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. The inactive precursors generally are activated when the prosequences are removed. Some or all of
30 the prosequences may be removed prior to activation. Such precursor protein are generally called proproteins.

Homologous sequences are found when there is an identity of sequence and can be determined upon comparison of sequence information, nucleic acid or amino acid, or through hybridization reactions between a known FtsZ and a candidate source. Conservative changes, such as Glu/Asp, Val/Ile, Ser/Thr, Arg/Lys and Gln/Asn can also be considered in determining sequence homology. Typically, a lengthy nucleic acid sequence can show as little as 50-60% sequence identity, and more preferably at least about 70% sequence identity, between the target sequence and the given FtsZ sequence of interest excluding any deletions which can be present, and still be considered related. Amino acid sequences are considered homologous by as little as 25% sequence identity between the two complete mature proteins. (See generally, Doolittle, R.F., *OF URFS and ORFS* (University Science Books, CA, 1986.)

In addition, not only can sequences provided herein be used to identify homologous FtsZ sequences, but the resulting sequences obtained therefrom can also provide a further method to obtain FtsZ sequences from other plant and/or bacterial sources. In particular, PCR can be a useful technique to obtain related FtsZ sequences from sequence data provided herein. One skilled in the art will be able to design oligonucleotide probes based upon sequence comparisons or regions of typically highly conserved sequence.

Once the nucleic acid sequence is obtained, the transcription, or transcription and translation (expression), of the FtsZ sequence in a host cell is desired to produce a ready source of the enzyme and/or modify the number and/or size of the plastids found therein. Other useful applications can be found when the host cell is a plant host cell, *in vitro* and *in vivo*.

Nucleic acids (genomic DNA, plasmid DNA, cDNA, synthetic DNA, mRNA, etc.) encoding FtsZ or amino acid sequences of the purified enzymes, which permit design of nucleic acid probes facilitating the isolation of DNA coding sequences therefor, are known in the art and are available for use in the methods of the present invention. It is generally recognized to an artisan skilled in the field to which the present invention pertains that the nucleic acid sequences provided herein and the amino acid sequences derived therefrom can be used to isolate other potential FtsZ genes from GenBank using DNA and peptide search techniques generally known in the art.

In addition to the sequences described in the present invention, DNA coding sequences useful in the present invention can be derived from algae, fungi, bacteria, plants, etc. Homology searches in existing databases using signature sequences corresponding to conserved nucleotide and amino acid sequences of FtsZ can be employed to isolate equivalent, related genes from other sources such as plants and microorganisms. Searches in EST databases can also be employed. Furthermore, the use of DNA sequences encoding enzymes functionally enzymatically equivalent to those disclosed herein, wherein such DNA sequences are degenerate equivalents of the nucleic acid sequences disclosed herein in accordance with the degeneracy of the genetic code, is also encompassed by the present invention. Demonstration of the functionality of coding sequences identified by any of these methods can be carried out by complementation of mutants of appropriate organisms, such as *E. coli*. The sequences of the DNA coding regions can be optimized by gene resynthesis, based on codon usage, for maximum expression in particular hosts.

The nucleic acid sequences which encode FtsZ can be used in various constructs, for example, as probes to obtain further sequences. Alternatively, these sequences can be used in conjunction with appropriate regulatory sequences to increase levels of the respective FtsZ sequence of interest in a host cell for recovery or study of the enzyme *in vitro* or *in vivo* or to decrease levels of the respective FtsZ sequence of interest for some applications when the host cell is a plant entity, including plant cells, plant parts (including but not limited to seeds, cuttings or tissues) and plants.

Thus, depending upon the intended use, the constructs can contain the nucleic acid sequence which encodes the entire FtsZ protein, or a portion thereof. For example, where antisense inhibition of a given FtsZ protein is desired, the entire FtsZ sequence is not required. Furthermore, where FtsZ constructs are intended for use as probes, it can be advantageous to prepare constructs containing only a particular portion of a FtsZ encoding sequence, for example a sequence which is discovered to encode a highly conserved FtsZ region.

As discussed above, nucleic acid sequence encoding a plant or other FtsZ proteins of this invention can include genomic, cDNA or mRNA sequence. By "encoding" is meant that the sequence corresponds to a particular amino acid sequence

either in a sense or anti-sense orientation. By "extrachromosomal" is meant that the sequence is outside of the plant genome of which it is naturally associated. By "recombinant" is meant that the sequence contains a genetically engineered modification through manipulation via mutagenesis, restriction enzymes, and the like.

5 A cDNA sequence may or may not contain pre-processing sequences, such as transit peptide sequences or targeting sequences to facilitate delivery of the FtsZ protein to a given organelle or membrane location. The use of any such precursor FtsZ DNA sequence is preferred for uses in plant cell expression. A genomic FtsZ sequence can contain the transcription and translation initiation regions, introns,
10 and/or transcript termination regions of the plant FtsZ, which sequences can be used in a variety of DNA constructs, with or without the FtsZ structural gene. Thus, nucleic acid sequences corresponding to the FtsZ sequences of this invention can also provide signal sequences useful to direct protein delivery into a particular organellar or membrane location, 5' upstream non-coding regulatory regions (promoters) having
15 useful tissue and timing profiles, 3' downstream non-coding regulatory regions useful as transcriptional and translational regulatory regions, and may lend insight into other features of the gene.

 Once the desired plant or other FtsZ nucleic acid sequence is obtained, it can be manipulated in a variety of ways. Where the sequence involves non-coding
20 flanking regions, the flanking regions can be subjected to resection, mutagenesis, etc. Thus, transitions, transversions, deletions, and insertions can be performed on the naturally occurring sequence. In addition, all or part of the sequence can be synthesized. In the structural gene, one or more codons can be modified to provide for a modified amino acid sequence, or one or more codon mutations can be
25 introduced to provide for a convenient restriction site or other purpose involved with construction or expression. The structural gene can be further modified by employing synthetic adapters, linkers to introduce one or more convenient restriction sites, or the like.

 For the most part, the constructs will involve regulatory regions functional in
30 plants which provide for altered size and number of plastids in a plant cell. The open reading frame, coding for the FtsZ protein, FtsZ-related protein or functional fragment thereof will be joined at its 5' end to a transcription initiation regulatory region such as

the wild-type sequence naturally found 5' upstream to the FtsZ or FtsZ-related structural gene, or to a heterologous regulatory region from a gene naturally expressed in plant tissues. Examples of useful plant regulatory gene regions include those from T-DNA genes, such as nopaline or octopine synthase, plant virus genes, such as
5 CaMV 35S, or from native plant genes.

The DNA sequence encoding a plant or other FtsZ protein of this invention can be employed in conjunction with all or part of the gene sequences normally associated with the FtsZ. In its component parts, a DNA sequence encoding FtsZ is combined in a DNA construct having, in the 5' to 3' direction of transcription, a
10 transcription initiation control region capable of promoting transcription and translation in a host cell, the DNA sequence encoding plant FtsZ and a transcription and translation termination region.

Potential host cells include both prokaryotic and eukaryotic cells. A host cell can be unicellular or found in a multicellular differentiated or undifferentiated organism
15 depending upon the intended use. Cells of this invention can be distinguished by having a FtsZ sequence foreign to the wild-type cell present therein, for example, by having a recombinant nucleic acid construct encoding a FtsZ protein therein not native to the host species.

Depending upon the host, the regulatory regions will vary, including regions
20 from viral, plasmid or chromosomal genes, or the like. For expression in prokaryotic or eukaryotic microorganisms, particularly unicellular hosts, a wide variety of constitutive or regulatable promoters can be employed. Expression in a microorganism can provide a ready source of the plant enzyme. Among transcriptional initiation regions which have been described are regions from bacterial
25 and yeast hosts, such as *E. coli*, *B. subtilis*, *Saccharomyces cerevisiae*, including genes such as beta-galactosidase, T7 polymerase, tryptophan E and the like.

In a preferred embodiment, the constructs will involve regulatory regions functional in plants which provide for modified production of plant FtsZ, and, possibly, modification of the plant cell plastid. The open reading frame coding for the
30 plant FtsZ or functional fragment thereof will be joined at its 5' end to a transcription initiation regulatory region. In embodiments wherein the expression of the FtsZ protein is desired in a plant host, the use of all or part of the complete plant FtsZ gene

is desired; namely all or part of the 5' upstream non-coding regions (promoter) together with the structural gene sequence and 3' downstream non-coding regions can be employed.

If a different promoter is desired, such as a promoter native to the plant host of interest or a modified promoter, i.e., having transcription initiation regions derived from one gene source and translation initiation regions derived from a different gene source, numerous transcription initiation regions are available which provide for a wide variety of constitutive or regulatable, e.g., inducible, transcription of the structural gene functions. The transcription/translation initiation regions corresponding to such structural genes are found immediately 5' upstream to the respective start codons. Among transcriptional initiation regions used for plants are such regions associated with the T-DNA structural genes such as for nopaline and mannopine synthases, the 19S and 35S promoters from CaMV, and the 5' upstream regions from other plant genes such as napin, ACP, SSU, PG, zein, phaseolin E, and the like. Enhanced promoters, such as double 35S, are also available for expression of FtsZ sequences. For such applications when 5' upstream non-coding regions are obtained from other genes regulated during seed maturation, those preferentially expressed in plant embryo tissue, such as ACP and napin-derived transcription initiation control regions, are desired. Such "seed-specific promoters" can be obtained and used in accordance with the teachings of issued U.S. Patent Numbers 5,608,152 and 5,530,194, which references are hereby incorporated by reference. Transcription initiation regions which are preferentially expressed in seed tissue, i.e., which are undetectable in other plant parts, are considered desirable for TAG modifications in order to minimize any disruptive or adverse effects of the gene product.

Regulatory transcript termination regions can be provided in DNA constructs of this invention as well. Transcript termination regions can be provided by the DNA sequence encoding the plant FtsZ or a convenient transcription termination region derived from a different gene source, for example, the transcript termination region which is naturally associated with the transcript initiation region. Where the transcript termination region is from a different gene source, it will contain at least about 0.25 kb, preferably about 1-3 kb of sequence 3' to the structural gene from which the termination region is derived.

Plant expression or transcription constructs having a plant FtsZ as the DNA sequence of interest for increased or decreased expression thereof can be employed with a wide variety of plant life, particularly, plant life involved in the production of vegetable oils for edible and industrial uses. Most especially preferred are temperate
5 oilseed crops. Plants of interest include, but are not limited to, rapeseed (Canola and High Erucic Acid varieties), sunflower, safflower, cotton, soybean, peanut, coconut and oil palms, and corn. Depending on the method for introducing the recombinant constructs into the host cell, other DNA sequences can be required. Importantly, this invention is applicable to dicotyledenous and monocotyledenous species alike and
10 will be readily applicable to new and/or improved transformation and regulation techniques.

The method of transformation is not critical to the instant invention; various methods of plant transformation are currently available. As newer methods are available to transform crops, they can be directly applied hereunder. For example,
15 many plant species naturally susceptible to *Agrobacterium* infection can be successfully transformed via tripartite or binary vector methods of *Agrobacterium*-mediated transformation. In addition, techniques of microinjection, DNA particle bombardment, and electroporation have been developed which allow for the transformation of various monocot and dicot plant species.

20 In developing the DNA construct, the various components of the construct or fragments thereof will normally be inserted into a convenient cloning vector which is capable of replication in a bacterial host, e.g., *E. coli*. Numerous vectors exist that have been described in the literature. After each cloning, the plasmid can be isolated and subjected to further manipulation, such as restriction, insertion of new fragments,
25 ligation, deletion, insertion, resection, etc., so as to tailor the components of the desired sequence. Once the construct has been completed, it can then be transferred to an appropriate vector for further manipulation in accordance with the manner of transformation of the host cell.

Normally, included with the DNA construct will be a structural gene having
30 the necessary regulatory regions for expression in a host and providing for selection of transformant cells. The gene can provide for resistance to a cytotoxic agent, e.g. antibiotic, heavy metal, toxin, etc., complementation providing prototrophy to an

auxotrophic host, viral immunity or the like. Depending upon the number of different host species in which the expression construct or components thereof are introduced, one or more markers can be employed, where different conditions for selection are used for the different hosts. A number of markers have been developed for use for
5 selection of transformed plant cells, such as those which provide resistance to various antibiotics, herbicides, or the like. The particular marker employed is not essential to this invention, one or another marker being preferred depending on the particular host and the manner of construction.

As mentioned above, the manner in which the DNA construct is introduced
10 into the plant host is not critical to this invention. Any method which provides for efficient transformation can be employed. Various methods for plant cell transformation include the use of Ti- or Ri-plasmids, microinjection, electroporation, DNA particle bombardment, liposome fusion, or the like. In many instances, it will be desirable to have the construct bordered on one or both sides by T-DNA,
15 particularly having the left and right borders, more particularly the right border. This is particularly useful when the construct uses *A. tumefaciens* or *A. rhizogenes* as a mode for transformation, although the T-DNA borders can find use with other modes of transformation.

Once a transgenic plant is obtained which contains cells with altered numbers
20 and/or sizes of chloroplasts, tissue containing such cells can then be used in plastid transformation experiments. For example, utilizing tissue containing cells with larger plastids provides for a larger target in plastid transformation methods, thus allowing for an increased probability of introduction of the foreign DNA into the plant cell plastid.

25 The DNA sequences, or polynucleotides, for use in plastid transformation of this invention will contain a plastid expression construct generally comprising a promoter functional in a plant cell plastid, and a DNA sequence of interest to be expressed in the transformed plastid cells.

Constructs and methods for use in transforming the plastids of higher plants
30 are described in Zoubenko *et al.* (*Nuc Acid Res* (1994) 22(19):3819-3824), Svab *et al.* (*Proc. Natl. Acad. Sci.*(1990) 87:8526-8530 and *Proc. Natl. Acad. Sci.*(1993) 90:913-917) and Staub *et al.* (*EMBO J.* (1993) 12:601-606). Constructs and methods for use

in transforming plastids of higher plants to express DNA sequences under the control of a nuclearly encoded, plastid targeted T7 polymerase are described in U.S. Patent Number 5,576,198. The complete DNA sequences of the plastid genome of tobacco are reported by Shinozaki *et al.* (*EMBO J.* (1986) 5:2043-2049).

5 Stable transformation of tobacco plastid genomes by particle bombardment is reported (Svab *et al.* (1990), *supra*) and Svab *et al.* (1993), *supra*). The methods described therein can be employed to obtain plants homoplasmic for plastid expression constructs using the methods described herein. Briefly, such methods involve DNA bombardment of a target host explant, preferably from a tissue which is
10 rich in metabolically active plastid organelles, such as green plant tissues including leaves, and cotyledons. The bombarded tissue is then cultured for ~2 days on a cell division promoting media. The plant tissue is then transferred to a selective media containing an inhibitory amount of the particular selective agent, as well as the particular hormones and other substances necessary to obtain regeneration for that
15 particular plant species. For example, in the above publications and the examples provided herein, the selective marker is the bacterial *aadA* gene and the selective agent is spectinomycin. The *aadA* gene product allows for continued growth and greening of cells whose chloroplasts comprise the marker gene product. Cells which do not contain the marker gene product are bleached. The bombarded explants will
20 form green shoots in approximately 3-8 weeks. Leaves from these shoots are then subcultured on the same selective media to ensure production and selection of homoplasmic shoots. As an alternative to a second round of shoot formation, the initial selected shoots can be grown to mature plants and segregation relied upon to provide transformed plants homoplasmic for the inserted gene construct.

25 The transformed plants so selected can then be analyzed to determine whether the entire plastid content of the plant has been transformed (homoplasmic transformants). Typically, following two rounds of shoot formation and spectinomycin selection, approximately 50% of the transgenic plantlets analyzed are homoplasmic as determined by Southern blot analysis of plastid DNA. These plantlets
30 are selected for further cultivation, both for analysis of the transgenic plastid phenotype (where the nuclear viral polymerase expression construct is also present in

the plastid transformant), or for use in methods to transform the viral polymerase construct into the nucleus of the transplastomic plants.

The methods of the present invention provide for a more efficient approach to obtaining homoplasmic plants. Wild-type plant cells typically contain 50 to 100 plastids per cell. However, once a transplastomic plant is obtained, the DNA sequence contained in the plant cell nucleus can be crossed away from the transplastomic cells. The DNA sequence transformed into the nucleus encoding for the alteration can be crossed away from the plant containing the transformed plastids. Once the DNA sequence has been crossed out, the plastids in the host plant cell can divide and revert back to normal (i.e. wild-type) plastid size and numbers. By applying the selective agent for which the plastid expression constructs provides resistance, cells containing a pure population of the plastids containing the foreign DNA can be obtained.

The vectors for use in plastid transformation preferably include means for providing a stable transfer of the plastid expression construct and selectable marker construct into the plastid genome. This is most conveniently provided by regions of homology to the target plastid genome. The regions of homology flank the construct to be transferred and provide for transfer to the plastid genome by homologous recombination, via a double crossover into the genome. The complete DNA sequence of the plastid genome of tobacco has been reported (Shinozaki *et al.*, *EMBO J.* (1986) 5:2043-2049). Complete DNA sequences of the plastid genomes from liverwort (Ohyama *et al.*, *Nature* (1986) 322:572-574) and rice (Hiratsuka *et al.*, *Mol. Gen. Genet.* (1989) 217:185-194), have also been reported.

Where the regions of homology are present in the inverted repeat regions of the plastid genome (known as IRA and IRB), two copies of the transgene are expected per transformed plastid. Where the regions of homology are present outside the inverted repeat regions of the plastid genome, one copy of the transgene is expected per transformed plastid. The regions of homology within the plastid genome are approximately 1kb in size. Smaller regions of homology can also be used, and as little as 100 bp can provide for homologous recombination into the plastid genome. However, the frequency of recombination and thus the frequency of obtaining plants having transformed plastids decreases with decreasing size of the homology regions.

Examples of constructs comprising such regions of homology for tobacco plastid transformation are described in Svab *et.al.* (1990 *supra*) and Svab and Maliga (1993 *supra*). Regions useful for recombination into tobacco and *Brassica* plastid genomes are also described in the following examples. Similar homologous recombination and selection constructs can be prepared using plastid DNA from the target plant species.

Other means of transfer to the plastid genome are also considered herein, such as by methods involving the use of transposable elements. For example, the constructs to be transferred into the plastid genome can be flanked by the inverted repeat regions from a transposable marker which functions in plant plastids. A DNA construct which provides for transient expression of the transposase required to transfer the target DNA into the plastids is also introduced into the chloroplasts. In this manner, a variety of phenotypes can be obtained in plants transformed with the same expression construct depending on positional effects which can result from insertion of the expression constructs into various locations on the plastid genome. Appropriate transposons for use in such plastic transformation methods include bacterial Tn10, bacteriophage Mu and various other known bacterial transposons.

The DNA sequence of interest in the plastid promoter expression constructs can be an encoding sequence which is oriented for expression of a particular structural gene, such that the protein encoded by the structural gene sequence is produced in the transformed plastid. In addition, the DNA sequence of interest can include a number of individual structural gene encoding regions such that an operon for expression of a number of genes from a single plastid promoter region is produced. Thus, it is possible to introduce and express multiple genes from an engineered or synthetic operon or from a pre-existing prokaryotic gene cluster. Such a method would allow large scale and inexpensive production of valuable proteins and fine chemicals in a particular desired plant tissue or a particular stage of development, depending upon the promoter used to drive nuclear expression of the specific viral polymerase. Such an approach is not practical by standard nuclear transformation methods since each gene must be engineered into a monocistron including an encoded transit peptide for plastid uptake and appropriate promoter and terminator signals. As a result, gene expression levels would be expected to vary widely between cistrons, and generation of a number of transgenic plant lines would be required. Ultimately crosses would be

required to introduce all of these cistrons into one plant to get expression to the target biochemical pathway.

Alternatively, the DNA sequence of interest in the plastid construct can be a fragment of an endogenous plastid gene oriented such that an RNA complementary to the endogenous gene mRNA is produced in the transformed plastid. Such antisense
5 constructs can be used to decrease the expression of the target plastid gene.

In order to provide a means of selecting the desired plant cells following plastid transformation, the polynucleotides for plastid transformation will also contain a construct which provides for expression of a marker gene. Expression of the marker
10 gene product allows for selection of plant cells comprising plastid organelles which are expressing the marker protein. In the examples provided herein, a bacterial *aadA* gene is expressed under the regulatory control of chloroplast 5' promoter and 3' transcription termination regions. The use of such an expression construct for plastid transformation of plant cells has been described by Svab and Maliga (1993, *supra*).
15 Expression of the *aadA* gene confers resistance to spectinomycin and streptomycin, and thus allows for the identification of plant cells expressing this marker gene. Selection for the *aadA* marker gene is based on identification of plant cells which are not bleached by the presence of streptomycin, or more preferably spectinomycin, in the plant growth medium. Other genes which encode a product involved in
20 chloroplast metabolism can also be used as selectable markers. For example, genes which provide resistance to plant herbicides such as glyphosate, bromoxynil or imidazolinone can find particular use. Such genes have been reported by Stalker *et al.* (*J. Biol. Chem.* (1985) 260:4724-4728; glyphosate resistant EPSP), Stalker *et al.* (*J. Biol. Chem.* (1985) 263:6310-6314; bromoxynil resistant nitrilase gene), and
25 Sathasivan *et al.* (*Nucl. Acids Res.* (1990) 18:2188; AHAS imidazolinone resistance gene).

The present invention also provides methods for obtaining a plastid transformed plant on medium containing glyphosate. At the initial event of transformation only a few plastids out of the many present in a plant cell are
30 transformed and therefore are able to express glyphosate resistant marker gene product. The rest of the untransformed plastids within the cell remains vulnerable to the effect of glyphosate. Therefore, although the cell contains transformed plastids, it

is unable to divide and sort out the transformed plastid resulting in lack of recovery of transformed callus tissue which would give rise to the transformed regenerants. Thus, any method that reduces plastid number to one or few within the cell has the potential to survive the effect of glyphosate and be useful as selectable marker for plastid

5 transformation.

The following examples are provided by way of illustration and not by way of limitation.

10

EXAMPLES

Example 1: Identification of Plant *ftsZ* Sequences

In order to obtain a plant tissue source with an altered number and/or size of
15 plastids using antisense and/or sense expression of the bacterial FtsZ plant
homologues, public as well as proprietary sequence databases are queried for
homologous sequences in soybean, rice, *Arabidopsis*, corn and *Brassica*. Two types
of plant FtsZ proteins have been previously identified in GenBank, type I FtsZ
proteins exemplified by accession gil1079731 (SEQ ID NO:32), appear to be imported
20 into the plastid, while type II FtsZ proteins, exemplified by accession gil3608494
(SEQ ID NO:33) and gil683524 (SEQ ID NO:34), appear to remain in the cytoplasm.
Homologs of both the type I FtsZ sequence as well as homologues of type II FtsZ
genes are described below. The sequences used to search against the databases are:
type I FtsZ homologue search was (SEQ ID NO:32), and for type II FtsZ searches,
25 (SEQ ID NO:33) is used.

Searches performed in proprietary databases containing sequences obtained
from *Arabidopsis* identified DNA sequences which are related to the FtsZ1 sequence.
The sequence of SEQ ID NO:1 is identified as AtFtsZ1. The deduced amino acid
sequence encoded by SEQ ID NO:1 is provided in SEQ ID NO:2. In addition, one
30 sequence (SEQ ID NO:3) was identified as related to the FtsZ2 sequence. The
deduced amino acid sequence encoded by SEQ ID NO:3 is provided in SEQ ID NO:4.

Sequences were also identified in databases containing sequences obtained from *Brassica*. One sequence was identified as related to the *Arabidopsis* FtsZ1 sequence. Based on sequence alignments between the two sequences, approximately 170 amino acids were predicted to be missing from the *Brassica* sequence at the N-terminus. To obtain a full length coding sequence for the *Brassica* FtsZ1 (BnFtsZ1) gene, RACE PCR using DNA obtained from *Brassica* leaves was performed using the primers SC258 (SEQ ID NO:35) and SC259 (SEQ ID NO:36). One reaction product was found to contain the most 5' sequence (SEQ ID NO:70) and was used to produce a full length sequence referred to as BnFtsZ1 (SEQ ID NO:5). The deduced amino acid sequence encoded by BnFtsZ1 is provided in SEQ ID NO:6)

A FtsZ1 homolog was also identified in tobacco with PCR using primers designed to the conserved amino acid domains of the *Arabidopsis* FtsZ1 sequence. The PCR primers used are identified as SC252 (SEQ ID NO:37), SC253 (SEQ ID NO:38), SC254 (SEQ ID NO:39) and SC255 (SEQ ID NO:40). The reaction products were cloned into TOPO TA (Invitrogen), and a single clone, referred to as xanthil-26-contig (SEQ ID NO:7), contained the most sequence. Additional primers were designed for use in RACE PCR to obtain full length coding sequence for the tobacco FtsZ1 homolog. For amplification of the 5' region, primers SC291 (SEQ ID NO:41) and SC292 (SEQ ID NO:42) were used, and for amplification of the 3' sequence, primers SC293 (SEQ ID NO:43) and SC294 (SEQ ID NO:44) were used. The PCR products were cloned in TOPO TA and sequenced. Clone xanftsZ1-5'-15 (SEQ ID NO:71) was chosen to be the best for the 5' tobacco FtsZ1 sequence since it contained the greatest amount of 5' sequence and overlap with xanthil-26-contig. This sequence was combined with the xanthil-26-contig to produce xanFtsZ1 (SEQ ID NO:8). The deduced amino acid sequence is provided in SEQ ID NO:9.

FtsZ homolog sequences were identified in databases containing DNA sequences obtained from corn by BLAST searches using the *Arabidopsis* FtsZ1 and FtsZ2 amino acid sequences. Ten sequence were identified as related to these FtsZ sequences, provided in SEQ ID NOs:10-19. The clones, when aligned, revealed six contigs, and the best representative clone for each were chosen for further analysis. Sequence analysis of SEQ ID NO:10 revealed a high homology to AtFtsZ1, and was estimated to be missing 158 amino acids at the N-terminal end when compared to

Arabidopsis FtsZ1. Clone SEQ ID NO:13 was found to overlap perfectly with SEQ ID NO:10 for 153 nt at the 5' end and in addition had 167 nt additional nt at the 5' end that had amino acid homology with the *Arabidopsis* FtsZ1. However, this clone was also not predicted to encode the full-length FtsZ, and was still missing 113 amino acids at the N-terminal end when compared to Arabidopsis FtsZ1. Interestingly, for clone SEQ ID NO:13, its homology with SEQ ID NO:10, ends at position 167nt and diverges. This could either be indicative of the presence of intronic sequence or a new class of FtsZ protein. Primer SC321 (SEQ ID NO:45) was designed to pull out the missing maize FtsZ1 sequence by RACE PCR.

Sequence analysis of SEQ ID NO:18 revealed its high homology to FtsZ2, and was also predicted to not to be full-length and missing about 286 amino acids at the N-terminal end when compared to Arabidopsis FtsZ2. Primer SC322 (SEQ ID NO:46) was designed to pull out the missing maize FtsZ2 sequence by RACE PCR. Although SEQ ID NO:14 and SEQ ID NO:15 were identified with the highest BLAST scores with FtsZ2.

Soybean FtsZ homolog sequences were identified in databases by BLAST searches with Arabidopsis FtsZ1 and FtsZ2 amino acid sequences. Twelve sequences were obtained, and are provided in SEQ ID NOs:20-31. Sequence analysis of SEQ ID NO:20, SEQ ID NO:24 and SEQ ID NO:25 revealed high homology to FtsZ1 and none to be full-length when compared to Arabidopsis FtsZ1. SEQ ID NO:25 had the longest sequence at the N-terminal end and is predicted to be missing 64 amino acids at the N-terminal when compared to Arabidopsis FtsZ1 sequence. Sequences of SEQ ID NO:20, SEQ ID NO:24 and SEQ ID NO:25 were used to correct the overlapping region. RACE PCR primers can now be designed to amplify the ends for obtaining a full length DNA sequence.

A sequence alignment between the Arabidopsis, Brassica, tobacco, soybean, and corn FtsZ1 protein sequences is provided in figure 1.

Example 2: Preparation of Plant Expression Constructs

2A. Nuclear Expression Constructs

Constructs are prepared for transformation into a plant cell nucleus for alteration of the plastid size and/or number in the transformed plant cell. Constructs can be prepared to alter the plastids constitutively, or in a tissue specific manner, for example, in leaf tissue, or seed tissue.

5 A plasmid containing the napin cassette derived from pCGN3223 (described in USPN 5,639,790, the entirety of which is incorporated herein by reference) was modified to make it more useful for cloning large DNA fragments containing multiple restriction sites, and to allow the cloning of multiple napin fusion genes into plant binary transformation vectors. An adapter comprised of the self annealed
10 oligonucleotide of sequence
CGCGATTTAAATGGCGCGCCCTGCAGGCGGCCGCTGCAGGGCGCGCCAT
TTAAAT (SEQ ID NO:47) was ligated into the cloning vector pBC SK+ (Stratagene) after digestion with the restriction endonuclease BssHII to construct vector pCGN7765. Plasmids pCGN3223 and pCGN7765 were digested with NotI and
15 ligated together. The resultant vector, pCGN7770, contains the pCGN7765 backbone with the napin seed specific expression cassette from pCGN3223.

The cloning cassette, pCGN7787, essentially the same regulatory elements as pCGN7770, with the exception of the napin regulatory regions of pCGN7770 have been replaced with the double CAMV 35S promoter and the tml polyadenylation and
20 transcriptional termination region.

A binary vector for plant transformation, pCGN5139, was constructed from pCGN1558 (McBride and Summerfelt, (1990) Plant Molecular Biology, 14:269-276). The polylinker of pCGN1558 was replaced as a HindIII/Asp718 fragment with a polylinker containing unique restriction endonuclease sites, AscI, PacI, XbaI, SmaI,
25 BamHI, and NotI. The Asp718 and HindIII restriction endonuclease sites are retained in pCGN5139.

A series of turbo binary vectors are constructed to allow for the rapid cloning of DNA sequences into binary vectors containing transcriptional initiation regions (promoters) and transcriptional termination regions.

30 The plasmid pCGN8618 was constructed by ligating oligonucleotides 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:48) and 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCC-3' (SEQ ID NO:49) into

SalI/XhoI-digested pCGN7770. A fragment containing the napin promoter, polylinker and napin 3' region was excised from pCGN8618 by digestion with Asp718I; the fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the napin promoter was closest to the blunted Asp718I site of pCGN5139 and the napin 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8622.

10 The plasmid pCGN8619 was constructed by ligating oligonucleotides 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCC -3' (SEQ ID NO:50) and 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:51) into SalI/XhoI-digested pCGN7770. A fragment containing the napin promoter, polylinker and napin 3' region was removed from pCGN8619 by digestion with Asp718I; the
15 fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the napin promoter was closest to the blunted Asp718I site of pCGN5139 and the napin 3' was closest to the blunted HindIII site was subjected to
20 sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8623.

The plasmid pCGN8620 was constructed by ligating oligonucleotides 5'-TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGGAGCT -3' (SEQ ID NO:52) and 5'-CCTGCAGGAAGCTTGCGGCCGCGGATCC-3' (SEQ ID NO:53) into
25 SalI/SacI-digested pCGN7787. A fragment containing the d35S promoter, polylinker and tml 3' region was removed from pCGN8620 by complete digestion with Asp718I and partial digestion with NotI. The fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow
30 fragment. A plasmid containing the insert oriented so that the d35S promoter was closest to the blunted Asp718I site of pCGN5139 and the tml 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert

orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8624.

The plasmid pCGN8621 was constructed by ligating oligonucleotides 5'-TCGACCTGCAGGAAGCTTGCGGCCGCGGATCCAGCT -3' (SEQ ID NO:54) and 5'-GGATCCGCGGCCGCAAGCTTCCTGCAGG-3' (SEQ ID NO:55) into 5 Sall/SacI-digested pCGN7787. A fragment containing the d35S promoter, polylinker and tml 3' region was removed from pCGN8621 by complete digestion with Asp718I and partial digestion with NotI. The fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested 10 with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the d35S promoter was closest to the blunted Asp718I site of pCGN5139 and the tml 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was 15 designated pCGN8625.

The plasmid construct pCGN8640 is a modification of pCGN8624 described above. A 938bp PstI fragment isolated from transposon Tn7 which encodes bacterial spectinomycin and streptomycin resistance (Fling et al. (1985), *Nucleic Acids Research* 13(19):7095-7106), a determinant for E. coli and Agrobacterium selection, 20 was blunt ended with Pfu polymerase. The blunt ended fragment was ligated into pCGN8624 that had been digested with SpeI and blunt ended with Pfu polymerase. The region containing the PstI fragment was sequenced to confirm both the insert orientation and the integrity of cloning junctions.

The spectinomycin resistance marker was introduced into pCGN8622 and 25 pCGN8623 as follows. A 7.7 Kbp AvrII-SnaBI fragment from pCGN8640 was ligated to a 10.9 Kbp AvrII-SnaBI fragment from pCGN8623 or pCGN8622, described above. The resulting plasmids were pCGN8641 and pCGN8643, respectively.

The Arabidopsis FtsZ1 nucleotide sequence was used to construct the sense 30 expression vector pCGN6495 for use in transformation of *Arabidopsis*, *Brassica* and tobacco. For this construct, the Arabidopsis ftsZ1 sequence was PCR amplified. To monitor protein expression of FtsZ1 in transformed lines, a c-myc tag (EQKLISEEDL

(SEQ ID NO:56)), was translationally fused to FtsZ1 at the C-terminal end. The PCR amplification was done by first round of amplification with primers SC247 (SEQ ID NO:57) and SC260 (SEQ ID NO:58) followed by amplification with SC247 (SEQ ID NO:59) and SC261 (SEQ ID NO:60) using the product of the first amplification as the
5 template DNA, using standard amplification parameters. The final amplification product, FtsZ1/c-myc fusion was cloned in the nuclear transformation vector pCGN8624 to create pCGN6495, which was used to nuclear transform Arabidopsis, canola and tobacco using standard protocols.

The turbo vector pCGN8624 was used for the antisense constructs such that
10 the antisense sequence is driven from d35S promoter. For *Arabidopsis* the coding sequence (from ATG to TAG) was amplified with primers SC248 (SEQ ID NO:61) and SC250 (SEQ ID NO:62) using AtFtsZ1 as template. For *Brassica*, primers SC276 (SEQ ID NO:63) and SC268 (SEQ ID NO:64) were used with PCR fragment SC3-1-1 (SEQ ID NO:70) as template DNA to generate a *HindIII/PstI* fragment and
15 cloned in pBSKS (Stratagene) to generate pCGN6528. Primer SC276 was designed to be located 140 bases downstream from ATG due to the presence of nonhomologous stretch of sequence compared to *Arabidopsis* FtsZ1 contained in the first 140 bases sequence fragment. The 3' half of the coding sequence was PCR amplified using primers SC269 (SEQ ID NO:65) and SC270 (SEQ ID NO:66) to produce a *PstI/NotI*
20 fragment, and subsequently cloned in pCGN6528 to generate pCGN6529. The *HindIII/PstI* fragment containing BnFtsZ1 sequence (from 140b downstream of ATG to TAG) was cloned in turbo vector pCGN8624 to generate final transformation vectors pCGN6530 and pCGN6611. The *HindIII/NotI* fragment containing BnFtsZ1 sequence was also cloned into pCGN8643 vector for seed-specific antisense FtsZ1
25 expression. For tobacco, primers SC305 and SC306 were designed to PCR amplify FtsZ1 sequence to produce a *SseI/NotI* fragment using 5' RACE PCR library DNA made from leaf RNA, and cloned into TOPO TA2.1 to produce pCGN6565. The *SseI/NotI* fragment from pCGN6565 was cloned in the turbo vector pCGN8624 to generate final transformation vector pCGN6566.

30

2A. Plastid Expression Constructs

Constructs and methods for use in transforming the plastids of higher plants are described in Zoubenko *et al.* (*Nuc Acid Res* (1994) 22(19):3819-3824), Svab *et al.* (*Proc. Natl. Acad. Sci.*(1990) 87:8526-8530 and *Proc. Natl. Acad. Sci.*(1993) 90:913-917) and Staub *et al.* (*EMBO J.* (1993) 12:601-606). Constucts and methods for use
5 in transforming plastids of higher plants to express DNA sequences under the control of a nuclearly encoded, plastid targeted T7 polymerase are described in U.S. Patent Number 5,576,198. The complete DNA sequences of the plastid genome of tobacco are reported by Shinozaki *et al.* (*EMBO J.* (1986) 5:2043-2049).

A plastid expression construct, pMON49218, was constructed to express the
10 synthetic CP4 sequence with the 14 amino acid GFP fusion from the promoter region of the 16SrDNA operon having the nuclear-encoded RNA polymerase region (PrmPEP+NEP), and the terminator region from the plastid rps16 gene. The DNA sequence of the Prm/NEP/G10L::14aaGFP fusion SEQ ID NO:67.

15

Example 3: Plant Transformation And Analysis

Constructs for the expression of sense or antisense sequences are transformed into tobacco cells using the methods described by Ursin *et. al.* (1991) *Plant Cell*
20 3:583-591.

Transgenic tobacco plants containing the nuclear FtsZ constructs were analyzed for alterations in plastid morphology, including size and number of plastids present in the plant cell.

Fifty-eight initial transformants (T1 generation) obtained from transformation
25 with FtsZ1 expression construct pCGN6495 were screened for the large plastid phenotype and divided into three categories. Thirty-four (34) lines contained less than 5 large plastids, 8 lines contained between 5-20 plastids and 16 lines more than 20 (wild-type# and more than wild-type#) plastids. One line, Nt6495-61, contained a single large plastid.

30 The screening method involved examining isolated mesophyll protoplasts at 100X magnification under light microscope. The large plastid containing transgenic

plants appear to be phenotypically indistinguishable from wild-type under culture and greenhouse conditions.

Estimation of plastid DNA copy number from several large plastid lines revealed no difference when compared to wild-type. Southern analysis was used to estimate transgene copy number in the large plastid lines and several lines with single integration events were identified. Western analysis of the large plastid lines with c-myc antibody confirmed expression of the introduced transgene (tagged by c-myc). T2 seeds were collected from selected plants from each of the three categories.

Example 4: Plastid Transformation and Analysis

Leaf material from three transgenic lines, Nt6495-30 (with <5plastids/cell), Nt6495-16 (with 5-20 plastids/cell) and Nt6495-69 (with 5-20 plastids/cell), were obtained for evaluation of plastid transformation efficiency and direct glyphosate selection. Plastid transformation vector pMON49218 which contains *aadA* gene for spectinomycin selection and GFP as a marker was used to bombard 15 leaf explants of each of the three transgenic lines. For each series of bombardment of the transgenic line 15 wild type control leaves were used. The order of bombardment for the transgenic line and the wild type leaves were randomized to eliminate any bias.

Transformation frequency of one event Nt6495-30 was approximately double that of the wild type control producing 7 versus 3 transformants respectively. Nt6495-16 and Nt6495-69 had approximately the same transformation frequency (3 transformants) as the control. Thus, our preliminary analysis reveals that plastid transformation efficiency can have been enhanced by reducing the plastid number from wild type to less than 5 plastids per cell. Interestingly, all of the plastid transformant regenerants from Nt6495 lines were very much slower in growth and size compared to those from wild type. It appears that the presence of the selectable antibiotic spectinomycin dihydrochloride at a concentration of 500mg/ml can have affected the regenerability of cells in the Nt6495 lines. Thus, it is possible that there could be more plastid transformed cells in the transgenic Nt6495 lines which were susceptible to the antibiotic and could not regenerate. To check if this was the case,

kill curves with lower concentrations of spectinomycin dihydrochloride (50, 100, 200, 300, 400 and 500mg/ml) can be used with each of the Nt6495-30, Nt6495-16 and Nt6495-69 lines to establish the concentration at which the regeneration of shoots are as good as in wild type. This concentration of spectinomycin dihydrochloride will then be used to repeat transformation frequency tests with the three Nt6495 lines.

To analyze for direct glyphosate selection, kill curves with varying levels of glyphosate will be established with the Nt6495 lines to find the best selection level. Plastid transformation vector pMON49218 will be used to bombard the Nt6495 lines and tested for direct selection using the optimized glyphosate level.

In Arabidopsis, FtsZ1 nuclear expression construct pCGN6495 was used to transform Columbia ecotype. T1 seeds were collected and about 100 kanamycin resistant seedlings were analyzed for alteration in plastid size and number following the same protocols as outlined for the tobacco section of this report. The transgenic plants were divided into three groups based on plastid number—I) 20 independent lines containing few large plastid (1-5), (II) 23 lines containing 5-20 plastids and (III) 50 lines containing wild-type plastid number were obtained. Selected T2 plants from each category were analyzed for number of transgene integration loci and sent to the growth chamber for T3 seed collection to identify homozygous plants. Such plants can be used in plastid transformations as described by Sikdar, *et al.* (1998) *Plant Cell Reports*, 18:20-24.

Transformed plants selected for *aadA* marker gene expression or glyphosate resistance are analyzed to determine whether the entire plastid content of the plant has been transformed (homoplasmic transformants). Typically, following two rounds of shoot formation and spectinomycin selection, approximately 50% of the transgenic plantlets which are analyzed are homoplasmic, as determined by Southern blot analysis of plastid DNA. Homoplasmic plantlets are selected for further cultivation.

Southern blot analysis is used to confirm the integration of the chimeric expression cassettes in the plastid genome. Preparation, electrophoresis, and transfer of DNA to filters is as described (Svab *et al.*, (1993 *supra*)). Total plant cellular DNA can be prepared as described by Dellaporta *et al.* (1983) *Plant Mol. Biol. Rep.* 1:19-21).

To visually observe the expression of marker genes such as GFP from the chloroplasts of transformed plants, various tissues are visualized utilizing a dissecting microscope. Protoplasts and chloroplasts are isolated as described in Sidorov, *et al.* (1994) *Theor. Appl. Genet.* 88:525-529.

5 The above results demonstrate that the sequences of the present invention provide an efficient means for the production of plastid transformed plants. Furthermore, such methods find use in plastid transformation methods involving the selection of transplastomic plants on herbicides, for example glyphosate.

10 All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

15 Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claim.

CLAIMS

What is claimed is:

- 5 1. In a method for transforming a plant cell plastid comprising the steps of introducing into cells of a plant a construct comprising a promoter functional in a plant cell plastid operably associated with a DNA sequence of interest and transforming said plant cell plastid with said construct,
 wherein the improvement comprises introducing said construct into a plant cell
10 having an altered plant plastid morphology selected from the group consisting of altered plastid size and altered plastid number in said plant cell.
2. The method according to Claim 1, wherein said plastid size is increased from a wild-type plant plastid morphology.
3. The method according to Claim 2, further wherein said plastid number is
15 decreased from a wild-type plant plastid morphology.
4. The method according to Claim 1, wherein said plastid size is decreased from a wild-type plant plastid morphology.
5. The method according to Claim 4, further wherein said plastid number is increased from a wild-type plant plastid morphology.
- 20 6. The method according to Claim 2, wherein said plant cell is obtained from a plant tissue source in which plastid division is inhibited.
7. The method according to Claim 6, wherein said plastid division is inhibited by introduction into cells of the plant tissue source a second DNA construct comprising in the 5' to 3' direction of transcription a promoter functional in a plant
25 cell, a DNA sequence coding for a gene involved in plastid cell division and a transcriptional termination sequence functional in a plant cell.
8. The method according to Claim 7, wherein said DNA sequence is in an antisense orientation.
9. The method according to Claim 8, wherein said construct contains a DNA
30 sequence coding for an FtsZ protein.
10. The method according to Claim 7, wherein said DNA sequence is in a sense orientation.

11. The method according to Claim 10, wherein said DNA sequence provides for sense suppression.

12. The method according to Claim 6, wherein said plastid division is inhibited by growing a plant under culture conditions which inhibit the division of plant cell plastids.

13. The method according to Claim 12, wherein said culture conditions comprise growing the plant tissue source under exposure to an inhibitor of bacterial cell division.

14. The method according to Claim 13, wherein said inhibitor is 5,5'-Bis-(8-anilino-1-naphthalenesulfonate).

15. The method according to Claim 6, wherein said plastid division is inhibited by genetic mutagenesis.

16. An isolated DNA sequence encoding a plant FtsZ protein from *Arabidopsis thaliana*.

17. The DNA sequence of Claim 16, wherein said FtsZ protein is encoded by a sequence which includes a sequence selected from the group consisting of SEQ ID Nos:1 and 3.

18. An isolated DNA sequence encoding a plant FtsZ protein from *Brassica*.

19. The DNA sequence of Claim 18, wherein said FtsZ protein is encoded by a sequence of SEQ ID NO:5.

20. An isolated DNA sequence encoding a plant FtsZ protein from soybean.

21. The DNA sequence of Claim 20, wherein said FtsZ protein is encoded by a sequence which includes a sequence selected from the group consisting of SEQ ID NOs:20-31.

22. An isolated DNA sequence encoding a plant FtsZ protein from corn.

23. The DNA encoding sequence of Claim 22, wherein said FtsZ protein is encoded by a sequence which includes a sequence selected from the group consisting of SEQ ID Nos:10-19.

24. A recombinant DNA construct comprising any of the DNA encoding sequences of Claims 16-23.

25. A plant cell comprising the DNA construct of 24.

26. A plant comprising a cell of Claim 25.

27. A method for improving the selectability of plant comprising,
transforming a plant cell source having an altered plastid morphology
with a construct comprising a promoter functional in a plant cell plastid operably
associated with a nucleic acid sequence encoding a selectable marker.
- 5 28. The method according to Claim 27, wherein said nucleic acid sequence
encodes an herbicide tolerance gene.
29. The method according to Claim 27, wherein said nucleic acid sequence
encodes a glyphosate tolerance gene.
30. A method for preparing a plant cell source with increased plastid
10 transformation efficiency comprising,
transforming a plant cell with a construct comprising a promoter functional in
plant cell operably associated with a nucleic acid sequence encoding a FtsZ protein.
31. A method for transforming a plant cell plastid comprising,
introducing into a plant cell having altered plastid morphology a first
15 nucleic acid construct comprising a promoter functional in a plant cell plastid
operably associated with a nucleic acid sequence of interest.

[illegible]

Figure 1

SEQUENCE LISTING

<110> Calgene LLC

<120> Methods for Transforming Plastids

<130> 15595/00/WO

<150> 60/109,892

<151> 1998-11-25

<160> 73

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1495

<212> DNA

<213> Arabidopsis sp

<400> 1

atggcgataa	ttccggttagc	acagcttaat	gagctaacga	tttcttcac	ttcttcttcg	60
tttcttacca	aatcgatata	ttctcattcg	ttgcatagta	gctgcatttg	cgcaagttct	120
agaatcagtc	aattccgtgg	cggcttctct	aaacgaagaa	gcgattcaac	aaggtctaag	180
tcgatcgcat	tgaggtgttc	cttctctccg	atggaatctg	cgagaattaa	ggtgattggt	240
gtcgggtggtg	gtggttaacaa	tgccgttaac	cggatgattt	caagcggttt	acagagtgtt	300
gatttctatg	cgataaacac	ggattcgcaa	gctctgttac	agtcttctgc	tgagaaccca	360
cttcaaattg	gagaactttt	aactcgtggg	cttggcactg	gtggaaaccc	gcttcttgga	420
gaacaagctg	cagaagaatc	aaaagatgca	attgctaata	ctcttaaagg	atcagacctt	480
gttttcataa	ctgctgggtat	gggtgggtgga	acaggggtctg	gtgctgcacc	tgtggtagct	540
cagatttcga	aggatgctgg	ttatttgact	gttgggtgtg	ttacctatcc	gtttagcttt	600
gaaggacgta	aaagatcttt	gcaggcactg	gaagctattg	aaaagctcca	aaagaatgtt	660
gataccctta	tcgtgattcc	aaatgatcgt	ctgctagata	ttgctgatga	acagacgcca	720
cttcaggacg	cgtttcttct	tgcagatgat	gttttacgcc	aaggagtaca	aggaatctca	780
gatattatta	ctataacctg	actagtcaat	gtagattttg	cagatgtgaa	ggcagtcag	840
aaagattctg	gaactgcaat	gctcggggta	ggtgtttctt	ccagcaaaaa	ccgggcagaa	900
gaagcagctg	aacaagcaac	tttggtctca	ttgatcggat	catccataca	atcagctact	960
ggtgtcgtct	acaacatcac	tgggtggaaaa	gacataactt	tgcaggaagt	gaaccgagta	1020
tcacaggctc	tgacaagtgt	ggcagaccca	tcggccaaca	tcataatttg	agctgttgtg	1080
gatgatcgct	acaccggaga	gattcatgta	acgataatcg	ccacaggctt	ctctcagtc	1140
ttccagaaga	cacttctgac	tgatccaaga	gcagctaaac	tccttgacaa	aatgggatca	1200
tcagggtcaac	aagagaacaa	aggaatgtct	ctgcctcacc	agaagcagtc	tccatcaact	1260
atctctacca	aatcgtcttc	tccccgtaga	cttttcttct	agttttcttt	ttttcctttt	1320
cggtttcaag	catcaaaaat	gtaacgatct	tcagggtcaa	atatcaatta	catttgattt	1380
tcaaaaaaaa	aaaaaaaaag	cggccgctct	agaggatcca	agcttacgta	cgcgtgcatg	1440
cgacgtcata	gctcttctat	agtgtcacct	aaattcaatt	cactggccgt	cgtttt	1495

<210> 2

<211> 433

<212> PRT

<213> Arabidopsis sp

<400> 2

Met Ala Ile Ile Pro Leu Ala Gln Leu Asn Glu Leu Thr Ile Ser Ser
 1 5 10 15
 Ser Ser Ser Ser Phe Leu Thr Lys Ser Ile Ser Ser His Ser Leu His
 20 25 30
 Ser Ser Cys Ile Cys Ala Ser Ser Arg Ile Ser Gln Phe Arg Gly Gly
 35 40 45
 Phe Ser Lys Arg Arg Ser Asp Ser Thr Arg Ser Lys Ser Met Arg Leu
 50 55 60
 Arg Cys Ser Phe Ser Pro Met Glu Ser Ala Arg Ile Lys Val Ile Gly
 65 70 75 80
 Val Gly Gly Gly Gly Asn Asn Ala Val Asn Arg Met Ile Ser Ser Gly
 85 90 95
 Leu Gln Ser Val Asp Phe Tyr Ala Ile Asn Thr Asp Ser Gln Ala Leu
 100 105 110
 Leu Gln Phe Ser Ala Glu Asn Pro Leu Gln Ile Gly Glu Leu Leu Thr
 115 120 125
 Arg Gly Leu Gly Thr Gly Gly Asn Pro Leu Leu Gly Glu Gln Ala Ala
 130 135 140
 Glu Glu Ser Lys Asp Ala Ile Ala Asn Ala Leu Lys Gly Ser Asp Leu
 145 150 155 160
 Val Phe Ile Thr Ala Gly Met Gly Gly Thr Gly Ser Gly Ala Ala
 165 170 175
 Pro Val Val Ala Gln Ile Ser Lys Asp Ala Gly Tyr Leu Thr Val Gly
 180 185 190
 Val Val Thr Tyr Pro Phe Ser Phe Glu Gly Arg Lys Arg Ser Leu Gln
 195 200 205
 Ala Leu Glu Ala Ile Glu Lys Leu Gln Lys Asn Val Asp Thr Leu Ile
 210 215 220
 Val Ile Pro Asn Asp Arg Leu Leu Asp Ile Ala Asp Glu Gln Thr Pro
 225 230 235 240
 Leu Gln Asp Ala Phe Leu Leu Ala Asp Asp Val Leu Arg Gln Gly Val
 245 250 255
 Gln Gly Ile Ser Asp Ile Ile Thr Ile Pro Gly Leu Val Asn Val Asp
 260 265 270
 Phe Ala Asp Val Lys Ala Val Met Lys Asp Ser Gly Thr Ala Met Leu
 275 280 285
 Gly Val Gly Val Ser Ser Ser Lys Asn Arg Ala Glu Glu Ala Ala Glu
 290 295 300
 Gln Ala Thr Leu Ala Pro Leu Ile Gly Ser Ser Ile Gln Ser Ala Thr
 305 310 315 320
 Gly Val Val Tyr Asn Ile Thr Gly Gly Lys Asp Ile Thr Leu Gln Glu
 325 330 335
 Val Asn Arg Val Ser Gln Val Val Thr Ser Leu Ala Asp Pro Ser Ala
 340 345 350
 Asn Ile Ile Phe Gly Ala Val Val Asp Asp Arg Tyr Thr Gly Glu Ile
 355 360 365
 His Val Thr Ile Ile Ala Thr Gly Phe Ser Gln Ser Phe Gln Lys Thr
 370 375 380
 Leu Leu Thr Asp Pro Arg Ala Ala Lys Leu Leu Asp Lys Met Gly Ser
 385 390 395 400
 Ser Gly Gln Gln Glu Asn Lys Gly Met Ser Leu Pro His Gln Lys Gln
 405 410 415
 Ser Pro Ser Thr Ile Ser Thr Lys Ser Ser Ser Pro Arg Arg Leu Phe
 420 425 430
 Phe

<210> 3
 <211> 1611
 <212> DNA
 <213> Arabidopsis sp

<400> 3
 tgttggtgccc gctcagaaat ctgaatcttc tccaatcaga aactctccac ggcattacca 60
 aagccaagct caagatcctt tcttgaacct tcacccggaa atatctatgc ttagaggtga 120
 agggactagt acaatagtca atccaagaaa ggaaacgtct tctggacctg ttgtcgagga 180
 ttttgaagag ccattctgctc cgagtaacta caatgaggcg aggattaagg ttattggtgt 240
 gggaggtggt ggatcaaattg ctgtgaatcg tatgatagag agtgaaatgt cagggtgtgga 300
 gttctggatt gtcaacactg atatccaggc tatgagaatg tctcctgttt tgcctgataa 360
 taggttacaa attggttaagg agttgactag ggggtttagggt gctggaggaa atccagaaat 420
 cggatatgaat gctgctagag agagcaaaga agttattgaa gaagctcttt atgggtcaga 480
 tatggtcttt gtcacagctg gaatgggagg tgggaactggc actggtgcag cccctgtaat 540
 tgcaggaaatt gccaaaggcga tgggtatatt gacagttgggt attgccacaa cgctttcttc 600
 gtttgagggt cgaagaagaa ctgttcaggc tcaagaaggg cttgcatctc tcagagacaa 660
 tgttgacact ctcatcgctc ttccaaatga caagttgctt acagctgtct ctcagtctac 720
 tccggttaaca gaagcatttta atctagctga tgatatactc cgtcaggggg ttcgtgggat 780
 atctgatatc attacgattc ctggtttgggt gaatgtggat tttgctgatg tgagagctat 840
 aatggcaaat gcggggtctt cattgatggg aataggaaact gcgacaggaa agagtcgggc 900
 aagagatgct gcgctaaatg caatccaatc ccctttgtta gatattggga ttgagagagc 960
 cactggaatt gtttggaaaca ttactggcgg aagtgacttg acattgtttg aggtaaatgc 1020
 tgctgcgga gtaatatatg atcttgtcga tccaactgcc aatcttatat tcggtgctgt 1080
 tgtagatcca gccctcagcg gtcaagtaag cataaccctg atagctacgg gtttcaaacg 1140
 acaagaagag ggagaaggac gaacagttca gatggtacaa gcagatgctg cgtcagttgg 1200
 agctacaaga agaccctctt ctctcttttag agaaagcggg tcagtggaga tcccagagtt 1260
 cttgaagaag aaaggcagct ctcgttatcc ccgagttctaa agcccaatct aatcactacc 1320
 ctgcacactg cagcaataac aaacgtgtgt gtactggtag tctggtactg ccttctggga 1380
 tacagcaaga tgtgttgatg tatgatcaag aatctgtgtg ggtgtgtata tgttctgtca 1440
 ctgctctctg tcgtgttctt gaatagggtt ttttagaaat cggagtttct ctctatgtca 1500
 cttccaaaac aaaaaaggag aagaagaatc acacttctcg aaccataaac atacttataa 1560
 gattatgaga gtttttagcag aaattattgt caaaaaaaaa aaaaaaaaaa a 1611

<210> 4
 <211> 397
 <212> PRT
 <213> Arabidopsis sp

<400> 4
 Met Leu Arg Gly Glu Gly Thr Ser Thr Ile Val Asn Pro Arg Lys Glu
 1 5 10 15
 Thr Ser Ser Gly Pro Val Val Glu Asp Phe Glu Glu Pro Ser Ala Pro
 20 25 30
 Ser Asn Tyr Asn Glu Ala Arg Ile Lys Val Ile Gly Val Gly Gly
 35 40 45
 Gly Ser Asn Ala Val Asn Arg Met Ile Glu Ser Glu Met Ser Gly Val
 50 55 60
 Glu Phe Trp Ile Val Asn Thr Asp Ile Gln Ala Met Arg Met Ser Pro
 65 70 75 80
 Val Leu Pro Asp Asn Arg Leu Gln Ile Gly Lys Glu Leu Thr Arg Gly
 85 90 95
 Leu Gly Ala Gly Gly Asn Pro Glu Ile Gly Met Asn Ala Ala Arg Glu

Ser Lys Glu Val Ile Glu Glu Ala Leu Tyr Gly Ser Asp Met Val Phe
 100 115 120 125
 Val Thr Ala Gly Met Gly Gly Gly Thr Gly Thr Gly Ala Ala Pro Val
 130 135 140
 Ile Ala Gly Ile Ala Lys Ala Met Gly Ile Leu Thr Val Gly Ile Ala
 145 150 155 160
 Thr Thr Pro Phe Ser Phe Glu Gly Arg Arg Thr Val Gln Ala Gln
 165 170 175
 Glu Gly Leu Ala Ser Leu Arg Asp Asn Val Asp Thr Leu Ile Val Ile
 180 185 190
 Pro Asn Asp Lys Leu Leu Thr Ala Val Ser Gln Ser Thr Pro Val Thr
 195 200 205
 Glu Ala Phe Asn Leu Ala Asp Asp Ile Leu Arg Gln Gly Val Arg Gly
 210 215 220
 Ile Ser Asp Ile Ile Thr Ile Pro Gly Leu Val Asn Val Asp Phe Ala
 225 230 235 240
 Asp Val Arg Ala Ile Met Ala Asn Ala Gly Ser Ser Leu Met Gly Ile
 245 250 255
 Gly Thr Ala Thr Gly Lys Ser Arg Ala Arg Asp Ala Ala Leu Asn Ala
 260 265 270
 Ile Gln Ser Pro Leu Leu Asp Ile Gly Ile Glu Arg Ala Thr Gly Ile
 275 280 285
 Val Trp Asn Ile Thr Gly Gly Ser Asp Leu Thr Leu Phe Glu Val Asn
 290 295 300
 Ala Ala Ala Glu Val Ile Tyr Asp Leu Val Asp Pro Thr Ala Asn Leu
 305 310 315 320
 Ile Phe Gly Ala Val Val Asp Pro Ala Leu Ser Gly Gln Val Ser Ile
 325 330 335
 Thr Leu Ile Ala Thr Gly Phe Lys Arg Gln Glu Glu Gly Glu Gly Arg
 340 345 350
 Thr Val Gln Met Val Gln Ala Asp Ala Ala Ser Val Gly Ala Thr Arg
 355 360 365
 Arg Pro Ser Ser Ser Phe Arg Glu Ser Gly Ser Val Glu Ile Pro Glu
 370 375 380
 Phe Leu Lys Lys Lys Gly Ser Ser Arg Tyr Pro Arg Val
 385 390 395

<210> 5

<211> 1450

<212> DNA

<213> Brassica sp

<400> 5

atggcgatta	gtccggttggc	acagcttaac	gagctaccag	tctcttcctc	gtttcttgcg	60
acatcccact	cgctgcacag	taccagaatc	agtggcggct	tctcaaaaca	aaggtttaag	120
caaacacggt	tgagatgctc	cttctctccg	atggagtctg	cgaggattaa	ggtgggttgg	180
gtcggcgggt	gtggtaacaa	tgccgtcaat	cgcattgatt	ccagcggctt	acagagtgtt	240
gatttctatg	cgataaacac	ggactctcaa	gctctcttgc	agtcttctgc	gcagaaccct	300
cttcaaattg	gagagctcct	aactcgtggc	cttgggactg	gtgggaaccc	gcttctagga	360
gaacaagctg	ctgaggaatc	taaagacgcg	attgctaata	ctcttaaagg	atctgacctt	420
gytttcatta	ctgctggtat	gggtggtggc	actggctccg	gtgctgctcc	tggtgttgct	480
cagatctcga	aagacgctgg	ttatttgacc	gttggtgttg	ttacctatcc	cttcagcttc	540
gaaggctcga	aaagatcttt	gcaggcactt	gaagccattg	aaaagctgca	gaagaacgtg	600
gataccctca	tcgtgatacc	aaatgatcgt	ctcctagata	ttgctgatga	acagacgcct	660

```

cttcaagacg cttttcttct cgcggtatgat gttttgcggc aaggagttca aggaatctct 720
gatattatta ctatacctgg actggtcaat gtagattttg cggatgtgaa gtcgggttatg 780
aaagattccg gaactgcat gctcggggtg ggtgtttctt caagcaagaa ccgagcagaa 840
gaagcagctg agcaagccac tttggctcca ttgatcggat catccattca atcagctact 900
ggtgtcgtct acaacatcac cggtggaaaa gacattactt tgcaggaagt gaaccgagta 960
tctcaggtgg tgacaagttt ggcagaccca tcggccaaca tcatatttgg agctgttgtg 1020
gatgatcgat acactggaga gattcatgta acgataatag ccacggggtt ctcacagtct 1080
ttccagaaga cacttctcag tgatccaaga gcagctaaac tactcgacaa aacgggatca 1140
tcaggtcaac aacaagagaa caaaggcagt caccagaggc agtctcctgc aactatcaac 1200
accaaactcat cttctccccg tagattgttc ttctagtatc ttttgTTTTT taagcatatt 1260
cctttatcaa aaatgtaacg atcttcaggc tcaaatatca attacttttc tccagattat 1320
ctcaaaagaa gtaatttgtt aaaccaaaaa aaaaaaaaaa gggcgggccgc tctagaggat 1380
ccaagcttac gtacgcgtgc atgcgacgtc atagctcttc tatagtgtca cctaaattca 1440
attcactggc                                     1450

```

<210> 6

<211> 411

<212> PRT

<213> Brassica sp

<220>

<221> VARIANT

<222> (1)...(411)

<223> Xaa = Any Amino Acid

<400> 6

```

Met Ala Ile Ser Pro Leu Ala Gln Leu Asn Glu Leu Pro Val Ser Ser
1          5          10
Ser Phe Leu Ala Thr Ser His Ser Leu His Ser Thr Arg Ile Ser Gly
20
Gly Phe Ser Lys Gln Arg Phe Lys Gln Thr Arg Leu Arg Cys Ser Phe
35          40          45
Ser Pro Met Glu Ser Ala Arg Ile Lys Val Val Gly Val Gly Gly Gly
50          55          60
Gly Asn Asn Ala Val Asn Arg Met Ile Ser Ser Gly Leu Gln Ser Val
65          70          75
Asp Phe Tyr Ala Ile Asn Thr Asp Ser Gln Ala Leu Leu Gln Ser Ser
85          90          95
Ala Gln Asn Pro Leu Gln Ile Gly Glu Leu Leu Thr Arg Gly Leu Gly
100          105          110
Thr Gly Gly Asn Pro Leu Leu Gly Glu Gln Ala Ala Glu Glu Ser Lys
115          120          125
Asp Ala Ile Ala Asn Ala Leu Lys Gly Ser Asp Leu Xaa Phe Ile Thr
130          135          140
Ala Gly Met Gly Gly Gly Thr Gly Ser Gly Ala Ala Pro Val Val Ala
145          150          155
Gln Ile Ser Lys Asp Ala Gly Tyr Leu Thr Val Gly Val Val Thr Tyr
165          170          175
Pro Phe Ser Phe Glu Gly Arg Lys Arg Ser Leu Gln Ala Leu Glu Ala
180          185          190
Ile Glu Lys Leu Gln Lys Asn Val Asp Thr Leu Ile Val Ile Pro Asn
195          200          205
Asp Arg Leu Leu Asp Ile Ala Asp Glu Gln Thr Pro Leu Gln Asp Ala
210          215          220
Phe Leu Leu Ala Asp Asp Val Leu Arg Gln Gly Val Gln Gly Ile Ser

```

225	Asp	Ile	Ile	Thr	Ile	Pro	Gly	Leu	Val	Asn	Val	Asp	Phe	Ala	Asp	Val
					245					250					255	
	Lys	Ser	Val	Met	Lys	Asp	Ser	Gly	Thr	Ala	Met	Leu	Gly	Val	Gly	Val
				260					265					270		
	Ser	Ser	Ser	Lys	Asn	Arg	Ala	Glu	Glu	Ala	Ala	Glu	Gln	Ala	Thr	Leu
				275				280					285			
	Ala	Pro	Leu	Ile	Gly	Ser	Ser	Ile	Gln	Ser	Ala	Thr	Gly	Val	Val	Tyr
		290					295					300				
	Asn	Ile	Thr	Gly	Gly	Lys	Asp	Ile	Thr	Leu	Gln	Glu	Val	Asn	Arg	Val
305						310					315				320	
	Ser	Gln	Val	Val	Thr	Ser	Leu	Ala	Asp	Pro	Ser	Ala	Asn	Ile	Ile	Phe
					325					330					335	
	Gly	Ala	Val	Val	Asp	Asp	Arg	Tyr	Thr	Gly	Glu	Ile	His	Val	Thr	Ile
				340					345					350		
	Ile	Ala	Thr	Gly	Phe	Ser	Gln	Ser	Phe	Gln	Lys	Thr	Leu	Leu	Ser	Asp
			355					360					365			
	Pro	Arg	Ala	Ala	Lys	Leu	Leu	Asp	Lys	Thr	Gly	Ser	Ser	Gly	Gln	Gln
		370					375					380				
	Gln	Glu	Asn	Lys	Gly	Ser	His	Gln	Arg	Gln	Ser	Pro	Ala	Thr	Ile	Asn
385						390					395				400	
	Thr	Lys	Ser	Ser	Ser	Pro	Arg	Arg	Leu	Phe	Phe					
				405						410						

<210> 7

<211> 1295

<212> DNA

<213> Nicotiana sp

<400> 7

tgccggttaac	cggatgattt	caagcgggtt	acagggtggt	gacttctatg	ctataaacac	60
ggatgctcaa	gcactgctac	agtctgctgc	tgaaaacccg	cttcaaattg	gagaacttct	120
gactcgtggg	cttggtactg	gtggtaatcc	tcttttaggg	gaacaggcag	tgaggagtc	180
gaaggaagcc	attgcaaatt	ctctaaaagg	ttcagatatg	gtgttcataa	cagcaggaat	240
gggtggaggt	acaggatctg	gtgctgctcc	tggtgtggct	caaataagcaa	aagaagcagg	300
ctatttgact	gttggtgttg	tcacataccc	attcagcttt	gaaggacgta	aaagatccgt	360
gcaggctctg	gaagcaattg	aaaaacttca	gaaaaatgta	gataccctta	tagtaattcc	420
aatgaccgtc	tgctagatat	tgctgatgag	cagacaccac	ttcaagatgc	ttttcttctt	480
gctgatgatg	tattacgccca	aggtgtccaa	ggaatttccg	atataattac	tatactgggg	540
cttgtaaattg	tggattttgc	cgatgtaaag	gtagtgatga	aagattcttg	aactgctatg	600
cttgaggattg	gggtttcatc	aagcaagaac	cgtgctgaag	aagcagccga	acaagcaact	660
cttgcccttc	ttaattggat	cgtccattca	atcgccactg	gggtagtatc	caccattcca	720
ggaggaaaag	accataactt	tgcagaaaag	gaatagggtg	tctcagggtg	ttacagtctg	780
gctgatccct	cccgctaaca	tcataatttg	tgctgttggtg	gatgagcgct	acaatggcga	840
aatacacgtg	accataattg	caactgggtt	tacccagctc	tttcagaaga	ctcttctctc	900
tgaccacga	ggtgcaaagc	ttgttgataa	aggcccagta	atccaagaaa	gcatggcatc	960
acctgttacc	ctgagggtcat	caacctcacc	ttcgacaaca	tcacgaacac	ctactcggag	1020
gctgttcttt	tagctccttt	atatagtttg	ttacggcttc	atttttctct	tttcttactt	1080
ttttcttttt	tactttcttt	gtattttacat	gttttgctga	ttggtgtttg	catttggtctg	1140
tagacatagt	gatgattctt	atcaagtgc	tcacattcat	actcgaaaaa	aaaaaaaaaa	1200
aaaaaaagta	ctctgcgttg	ttaccctactg	ttaagggcga	attctgcaga	tatcccatca	1260
cactggcggc	cgctcgagca	tgcattctaga	gggcc			1295

<210> 8

<211> 1255

<212> DNA

<213> Nicotiana sp

<400> 8

atggccacca	tctcaaacc	agcagagata	gcagcttctt	ctccttcctt	tgctttttac	60
cactcttcct	ttattcctaa	acaatgctgc	ttcaccaaag	ctcgccggaa	aagcttatgt	120
aaacctcaac	gtttcagcat	ttcaagtcca	tttactcctt	ttgattctgc	taagattaag	180
gttatcggcg	tcggtggcgg	tggttaacaat	gccgttaacc	ggatgatttc	aagcgggtta	240
caggggtgttg	acttctatgc	tataaacacg	gatgctcaag	cactgctgca	gtctgctgct	300
gaaaacccgc	ttcaaattgg	agaacttctg	actcgtgggc	ttggtactgg	tggtaatcct	360
cttttagggg	aacaggcagc	ggaggagtcg	aaggaagcca	ttgcaaattc	tctaaaaggt	420
tcagatatgg	tggtcataac	agcaggaatg	ggtggaggta	caggatctgg	tgctgctcct	480
gttgtggctc	aaatagcaaa	agaagcaggc	tatttgactg	ttggtgttgt	cacataccca	540
ttcagctttg	aaggacgtaa	aagatccgtg	caggctctgg	aagcaattga	aaaacttcag	600
aaaaatgtag	atacccttat	agtaattccc	aatgaccgtc	tgctagatat	tgctgatgag	660
cagacaccac	ttcaagatgc	ttttcttctt	gctgatgatg	tattacgcca	aggtgtccaa	720
ggaatttccg	atataattac	tatacctggg	cttgtaaagt	tggtatttgc	cgatgtaaag	780
gtagtgatga	aagattctgg	aactgctatg	cttggaagtg	gggtttcatc	aagcaagaac	840
cgtgctgaag	aagcagccga	acaagcaact	cttgcccctc	ttattggatc	gtccattcaa	900
tcagccactg	gggtagtatc	caccattcca	ggaggaaaag	acataacttt	gcagaaagtg	960
aataggggtgt	ctcaggttgt	tacagtctgg	ctgatccctc	ccgctaacat	catatttggt	1020
gctgttggtg	atgagcgcta	caatggcgaa	atacacgtga	ccataattgc	aactggtttt	1080
acccagtctt	ttcagaagac	tcttctctct	gacccacgag	gtgcaaagct	tggtgataaa	1140
ggcccagtaa	tccaagaaag	catggcatca	cctgttaccc	tgaggtcatc	aacctcacct	1200
tcgacaacat	cacgaacacc	tactcggagg	ctgttctttt	agctccttta	tatag	1255

<210> 9

<211> 413

<212> PRT

<213> Nicotiana sp

<400> 9

Met	Ala	Thr	Ile	Ser	Asn	Pro	Ala	Glu	Ile	Ala	Ala	Ser	Ser	Pro	Ser
1				5					10					15	
Phe	Ala	Phe	Tyr	His	Ser	Ser	Phe	Ile	Pro	Lys	Gln	Cys	Cys	Phe	Thr
			20					25					30		
Lys	Ala	Arg	Arg	Lys	Ser	Leu	Cys	Lys	Pro	Gln	Arg	Phe	Ser	Ile	Ser
		35					40					45			
Ser	Ser	Phe	Thr	Pro	Phe	Asp	Ser	Ala	Lys	Ile	Lys	Val	Ile	Gly	Val
	50				55					60					
Gly	Gly	Gly	Gly	Asn	Asn	Ala	Val	Asn	Arg	Met	Ile	Ser	Ser	Gly	Leu
65				70				75						80	
Gln	Gly	Val	Asp	Phe	Tyr	Ala	Ile	Asn	Thr	Asp	Ala	Gln	Ala	Leu	Leu
			85					90						95	
Gln	Ser	Ala	Ala	Glu	Asn	Pro	Leu	Gln	Ile	Gly	Glu	Leu	Leu	Thr	Arg
		100						105					110		
Gly	Leu	Gly	Thr	Gly	Gly	Asn	Pro	Leu	Leu	Gly	Glu	Gln	Ala	Ala	Glu
		115					120					125			
Glu	Ser	Lys	Glu	Ala	Ile	Ala	Asn	Ser	Leu	Lys	Gly	Ser	Asp	Met	Val
	130					135					140				
Phe	Ile	Thr	Ala	Gly	Met	Gly	Gly	Gly	Thr	Gly	Ser	Gly	Ala	Ala	Pro
145					150					155				160	
Val	Val	Ala	Gln	Ile	Ala	Lys	Glu	Ala	Gly	Tyr	Leu	Thr	Val	Gly	Val
			165					170						175	
Val	Thr	Tyr	Pro	Phe	Ser	Phe	Glu	Gly	Arg	Lys	Arg	Ser	Val	Gln	Ala


```

      180      185      190
Leu Glu Ala Ile Glu Lys Leu Gln Lys Asn Val Asp Thr Leu Ile Val
      195      200      205
Ile Pro Asn Asp Arg Leu Leu Asp Ile Ala Asp Glu Gln Thr Pro Leu
      210      215      220
Gln Asp Ala Phe Leu Leu Ala Asp Asp Val Leu Arg Gln Gly Val Gln
      225      230      235
Gly Ile Ser Asp Ile Thr Ile Pro Gly Leu Val Asn Val Asp Phe
      245      250      255
Ala Asp Val Lys Val Val Met Lys Asp Ser Gly Thr Ala Met Leu Gly
      260      265      270
Val Gly Val Ser Ser Ser Lys Asn Arg Ala Glu Glu Ala Ala Glu Gln
      275      280      285
Ala Thr Leu Ala Pro Leu Ile Gly Ser Ser Ile Gln Ser Ala Thr Gly
      290      295      300
Val Val Ser Thr Ile Pro Gly Gly Lys Asp Ile Thr Leu Gln Lys Val
      305      310      315
Asn Arg Val Ser Gln Val Val Thr Val Trp Leu Ile Pro Pro Ala Asn
      325      330      335
Ile Ile Phe Gly Ala Val Val Asp Glu Arg Tyr Asn Gly Glu Ile His
      340      345      350
Val Thr Ile Ile Ala Thr Gly Phe Thr Gln Ser Phe Gln Lys Thr Leu
      355      360      365
Leu Ser Asp Pro Arg Gly Ala Lys Leu Val Asp Lys Gly Pro Val Ile
      370      375      380
Gln Glu Ser Met Ala Ser Pro Val Thr Leu Arg Ser Ser Thr Ser Pro
      385      390      395
Ser Thr Thr Ser Arg Thr Pro Thr Arg Arg Leu Phe Phe
      405      410

```

<210> 10
 <211> 1278
 <212> DNA
 <213> Zea mays

<220>
 <221> misc_feature
 <222> (1)...(1278)
 <223> n = A,T,C or G

```

<400> 10
gatcttgtct tcataacagc tgggatggga gggggtactg gatctggtgc tgctccagtt      60
gttgcccaga tatcaaagga agctggttat cttactgttg gtgttggtcac ctatccattc      120
agtttgcagg gccgtaagcg ctctgtacag gcattggaag cactagagaa gctggaaaag      180
agtgtagaca cacttattgt gattccaaat gataagttat tagatgttgc ggatgaaaac      240
atgcccttgc aagatgcatt tctccttgca gatgatgtcc ttcgtcaggg tgttcaagga      300
atatcagaca tcatcacaaat accgggactt gtcaatgttg attttgctga tgtaaaagct      360
gtcatgaaaa actctggaac tgccatgctc ggtgttggtg tttcttcag caaaaatcgg      420
gcccaagaag ctgctgaaca ggcaacactt gctccttga ttggatcatc catcgaggca      480
gctactggcg ttgtgtataa tattactggt gggaaggaca tcactttgca agaagtgaac      540
aaggtgtccc agattgtgac aagcctagct gacccatctg cgaacataat tttcgggtgct      600
gtcgttgatg accgttacac tggtagata catgtgacaa tcattgcgac aggatttcca      660
cagtccttcc agaaatccct tttggcggat ccaaaggag cacgtatagt ggaatccaaa      720
gagaaagcag caaccctcgc ccataaagca gcagcagctg cagttcaacc ggtccctgct      780
tctgcttggt ctcgaagact cttctcctga gaagctcatt tggatgaaccg tgactcgtag      840

```

tgcatatgat	ttgcatttag	cgtgttgagg	gcagtcctta	aggatgatctt	cggatatctg	900
gagatttata	gcttgggcta	gtgttcggta	gtggtagaat	aagtttcagt	gtatgtatcg	960
ttgctttgct	ttatgttttt	gaggatcagg	cgggtgaggct	gagagaagtg	ctcagcaact	1020
caacattgaa	ctgttgtaga	agatccttga	ttgcttttat	tgctgcaaca	tgccaacaac	1080
cctctgttgg	attcamcmna	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	1140
aaaaaaaaaa	aaaaaaaaaa	aaaanncaaa	aaaaaaaaaa	aaaaaaaaag	gcggccgccc	1200
actagtgagc	tcgtcgaccc	gggaattaat	tccggaccgg	tacctgcagg	cgtaccagct	1260
ttccctatag	tgagtcgt					1278

<210> 11
 <211> 283
 <212> DNA
 <213> Zea mays

<400> 11						60
gctccagttg	ttgccagat	atcaaaggaa	gctgggttatc	ttactgttgg	tggtgtcacc	120
tatccattca	gtttcgaggg	ccgtaagcgc	tctgtacagg	cattggaagc	actagagaag	180
ctggaaaaga	gtgtagacac	acttattgtg	attccaaatg	ataagttatt	agatgttgcg	240
gatgaaaaca	tgcccttgca	agatgcattt	ctccttgtag	atgatgtcct	tcgtcagggg	283
gttcaaggaa	tatcagacat	catcacaata	ccgggacttg	tca		

<210> 12
 <211> 287
 <212> DNA
 <213> Zea mays

<220>
 <221> misc_feature
 <222> (1)...(287)
 <223> n = A,T,C or G

<400> 12						60
ggggccgtaag	cgctctgtac	aggcattgga	agcactagag	aagctggaaa	agagtgtaga	120
cacacttatt	gtgattccan	atnatnngtt	attagatgtt	gcggatgaaa	acatgccctt	180
gcaagatgca	tttctccttg	cagatgatgt	ccttcgctag	ggtgttcaag	gaatatcaga	240
catcatcaca	ataccgggac	ttgtcaatgt	tgattttgct	gatgtaaaag	ctgtcatgaa	287
aaactctgga	actgccatgc	tcgggtgttg	tgttttcttc	agcaaaa		

<210> 13
 <211> 1122
 <212> DNA
 <213> Zea mays

<400> 13						60
gctataaaca	ccgattccca	agcccttatt	aattcacaag	cgcaatatcc	tctgcaaatt	120
ggagagcagt	tgaccgcggy	cttaggtgcc	ggtggaaatc	cgaatttggg	agagcaggct	180
gctgaggaat	caagagaaac	catagccact	gccctgaggg	attcagatct	tgtcttcata	240
acagctggga	tgaggagggg	tactggatct	ggtgctgctc	cagttgttgc	ccagatatca	300
aagggaagctg	gttatcttac	tggtgggtgtt	gtcacctatc	cattcagttt	cgagggccgt	360
aagcgctctg	tacaggcaaa	gtatctgagc	cccccttcac	tcctgaattt	taattcaaac	420
tgatcatatct	cgttctgcga	ctttcttttg	ctcgatggaa	gcattagttt	gtagtcataa	480
caatgacatc	cagccacatt	tattgctgat	gatgtataca	atggtaggtc	aaagaaatgt	540
agcatcatgc	catcacctgt	agttcatctc	atcattttgt	tcctactttt	ctgcgtgggt	600
gatgcccaaa	acaatatata	actatgtggt	tgatctgttg	cattgccttg	tggagggatg	660
tttatgttgt	gaaatatctc	aaaacacatg	tcattatgaa	tattccctcc	tggtggttgtg	

gggacttggt	tcaaagtcta	tgaattaaga	acaaggcaac	ataaagtgtt	aaatgttaac	720
cgtctttcgt	ccatgaaaca	ttattccctt	gaggataatg	ggccttggac	aaaggctgat	780
gagagtataa	ttaccaagct	taaatcttcg	taataaaaatt	tcaatagata	ttgtaagata	840
acataaaaata	aaggggtataa	aaaggggtaa	ataaatcata	gacgaattat	attatattta	900
cttaatatat	tgaatcattg	aatacaataa	tacctctgcc	ttggcaaagg	ttggattccg	960
aaaaatgtga	ttgcaagtta	ccagaatgcg	tgaacagtaa	aggaataactg	ttcactatct	1020
ataggcacag	gacacagcct	gtggaggaat	tcaattatac	ccgtcataag	agtttacaca	1080
ttgacttaga	cctttatgga	ctaaaagatc	attgctatct	tt		1122

<210> 14

<211> 291

<212> DNA

<213> Zea mays

<220>

<221> misc_feature

<222> (1)...(291)

<223> n = A,T,C or G

<400> 14

aaaatagtgt	ggacacccta	atcgatcatc	caaagtataa	gttgctgtct	gctgtttctc	60
caaatacacc	tgtaactgaa	gcattttaatc	tggtgatga	tattcttcgt	caaggcattc	120
gtggcatatc	tgatataatt	acggttccctg	ggnagggttaa	tggtgatttt	gctgacgtac	180
gtgctatcat	gcaaaatgca	gggtcatcct	tgatgggtat	agggactgct	acaggaaagt	240
caagagcaag	ggatgctgct	cttaacgcca	tccagtcgcc	gctgcttgat	a	291

<210> 15

<211> 415

<212> DNA

<213> Zea mays

<220>

<221> misc_feature

<222> (1)...(415)

<223> n = A,T,C or G

<400> 15

gagcaaggga	tgctgctctt	aacgccatcc	agtcgccgct	gcttgatatt	ggaattgaaa	60
gagccacagg	cattgtgttg	aatatcactg	ggggaactga	cctgactttg	tttgagggtga	120
atgctgcggc	cgaaattatc	tacgaccttg	tcgatccaaa	cgctaactctg	atattttggcg	180
ccgtcataga	cccgtcactg	agtgggcagg	tgagcataac	cttgatagct	actggcttca	240
aacggcagga	tgaaccagaa	ggccgcgtgt	cgaagggttg	gcaacaaggt	gagaatggcc	300
gacgcccac	cccagcanag	ggcaacaaca	cggtggaaat	tccaaaattc	ccgccaacaa	360
aaagggccct	tcnnccttc	cacnattttg	actggtcctg	tctgcacctg	tatga	415

<210> 16

<211> 744

<212> DNA

<213> Zea mays

<220>

<221> misc_feature

<222> (1)...(744)

<223> n = A,T,C or G

```

<400> 16
aattcccggg tgcacccacg cgtcccgcgg acgcgtgggt ggaatatcac tggagggaac      60
gatctaacct tgacagaggt gaatgctgca gctgaagtaa tctatgatct tgttgacctt      120
ggtgcaaatc tgatttttgg ctctgttata gatccgtcat acactgggtca agtgagcata      180
actctaattg caactgggtt caaacgccag gaggaaagtg agagccggtc ttcacaggct      240
ggaggagaca agcaaccgcg gtcgctcggc tggttttctc ccacttcca ggaggagggt      300
catgcattgc aaatccana gttcctacag aggaaagggc gtccagggtt tcacgagtct      360
gaacacactt tggatcaatg tttttcttgt catagtgttg tacgatgcag gtttggtttc      420
tggtgtctct aggtagcaag gtagaacaga tgttcctgaa cccgcacata ctaatctgtg      480
tgcaaacttc ngccgctgag taccattggc ttgggctgct ttgcttctca ngaacctgca      540
gtgagggtct aatttgctag ttagtatgat taaaagttaa gcgctgagac caaattatac      600
gttccgtgtg aatgattact tgctcncgtc cattttcttt tcaaaaaaaaa aaaaaaaaaa      660
aaaaaggcgg cgctntanag gatccaagct tacttcccc gcatncgacn canagctntt      720
ntatagngtg acctaaattc aatc                                     744

```

```

<210> 17
<211> 230
<212> DNA
<213> Zea mays

```

```

<220>
<221> misc_feature
<222> (1) ... (230)
<223> n = A,T,C or G

```

```

<400> 17
ggctgctgag gaatcaagag aaaccatagc cactgccctg agggattcag atcttgtttt      60
cataacagct gggatggnag ggggtgctgc tccaattggt gccagatat caaaggaagc      120
tggttatctt actgttggtg ttgtcaccta tccattcaat ttcgagggcc gtaagcgctc      180
tttacaggca agtatctgag ccccccttca ctctgaatt agaattcaaa      230

```

```

<210> 18
<211> 318
<212> DNA
<213> Zea mays

```

```

<220>
<221> misc_feature
<222> (1) ... (318)
<223> n = A,T,C or G

```

```

<400> 18
caggcattgt gtggaatatc actgggggaa ctgacctaac tttgtttgag gtgaatgctg      60
cggccgaaat tatctacgac cttgtcgatc caaatgctaa tctgatattt ggtgccgtca      120
tagaccgctc actgagtggg caggtgagca taacctgata gctactggct tcaaacggca      180
ggatgaacca gaaggccgcg tgtcgaaggg tgggcaacaa agtgagaatg gccgacgcc      240
gtcccccgca gagggcagca gcacgggtga gttccagagt cctgcgacgt agagganctt      300
ctcgcttccc agagttga                                     318

```

```

<210> 19
<211> 471
<212> DNA
<213> Zea mays

```

```

<220>

```

<221> misc_feature
 <222> (1)...(471)
 <223> n = A,T,C or G

<400> 19
 cgacgcccac ggtgacgaat gctgtcagcc acgctgtgct acacggggga aacaatgcaa 60
 anacattacc tgccctcactc ntgcttgctc ctgtaaatat aatgatngtc gctgctacat 120
 natatttact cctgctgctg cttgaggcca ttattctgta cgtaaatgaa gccactacta 180
 ctctcacaca gcatgcgccc gccgacgacg tacgtacgtg tattatatac gctctacccc 240
 gtgagctttt gttcgagtga tacgtgatcc atccatgcat ggatgcttat gtatgtatat 300
 gtgttagtcg tctcagggaa cggggcanca naaggggggtg ttgtattana ttacgtctt 360
 ctggtgatta aataanaaaag gggatatgtg gatgtgtgca aaaaaaaaaa aaaaanaaaa 420
 aaaaaaaaaa aaaaaaaaaa ggcggccgcc gactagttag ctcgtcgacc c 471

<210> 20
 <211> 1085
 <212> DNA
 <213> Glycine sp

<220>
 <221> misc_feature
 <222> (1)...(1085)
 <223> n = A,T,C or G

<400> 20
 cggctcgagg aaccctatta aaattggaga agttctgact cgtggattag gtacgggagg 60
 gaatccactt ttgggggaac aagctgcaga ggaatcaaga gatgctattg ctgatgctct 120
 taaaggatca gatttgggtg ttataacggc tgggatgggt gggggaaccg ggtctgggtg 180
 tgccccagtt gtagcccaaa tatcaaaaga ggcaggttac ttgactgtag gtgttggttac 240
 ctatcccttc agttttgaag gacgtaagag atccttgacg gcctttgaag ccacgaaaag 300
 gctgcagaaa aatggttgaca cmnttatagt gawtccmaat gmccgtctgc ttgacawagy 360
 tratragcar atgcctcttc aaggatgctt tccgytttgc agatgacgtt ytmaggcaag 420
 gagtmcaggg aatatcagac attatamctg tacctggact tkkcaaatgt ggattttgca 480
 agatgtaaaa gctgtgatga aagactctgg gactgcaatg cttggagtag gtgtttccty 540
 cggtaaaaaa ccgagcagaa gaagcagccg aacaggctac tttggctcct ttaattggat 600
 cctctattca gtcaagctac tggggtagtg tataatatta ctggagggaa aggacataac 660
 cctgcaggaa gtgracaggg tttymaggt kgkgacyark ttggctgatc cttctgctaa 720
 tattatatatt ggggctgtcg ttgatgatcg ctacacgggg gagattcacg tgactatcat 780
 tgcaactggc ttctcacagt cttttcagaa gaagttgcta acagatccaa gggcttgcaa 840
 agctgcttga caaggtggct gagggccaag aaagcaaggc agtccctcct cccctcaagt 900
 cctcaaacaa gggtgaatct agaccatccc cgcgaaagct ctttttttag ttgcatgggt 960
 ctttttacc tttttcattt ttccaattat tattattata ttatatnggc cgatcaaaaa 1020
 aaaaaaaaaa ggcggccgcc gactagttag ctcgtcgacc cgggaattaa ttccggaccg 1080
 gtacc 1085

<210> 21
 <211> 797
 <212> DNA
 <213> Glycine sp

<400> 21
 ccagctggcg aaaggggatg tgctgcaagg cgattaagtt gggtagcgag ggttttccca 60
 gtcacgacgt tgtaaaacga cggcagtgaa ttgaatttag gtgacactat agaagagcta 120
 tgacgtcgca tgcacgcgta cgtaagctcg gaattcggct cgagaggcta ctttggtctc 180
 ttttaattga tcctctattc agtcagctac tggggtagtg tataatatta ctggaggaaa 240

ggacataacc	ctgcaggaag	tgaacagggt	ttctcaggtt	gtgactagtt	tggtgatcc	300
ttctgctaatt	attatatattg	gggctgtcgt	tgatgatcgc	tacactgggg	agattcacgt	360
gactatcatt	gcaactggct	tctcacagtc	ttttcagaag	aagttgctaa	cagatccaag	420
ggctgcaaaag	ctgcttgaca	aggtggctga	gggccaaagaa	agcaaggtag	tccctcctcc	480
cctcaagtc	tcaaacaagg	ttgaatctag	accatccccg	cgaaagctct	tttttttagt	540
tgcatgggtc	tttttaccct	ttttcatatt	tccaattatt	attattatat	tattattggcc	600
gatcaaaaaa	aaaattatta	tattatatgg	taggacacaa	tgatcttgat	gcttaattaa	660
gtgagatatt	attctcttga	tgtaaaaaa	aaaaaaaaag	ggcgccgcgc	gactagtgag	720
ctcgtcgacc	cggaatttaa	ttccggaccg	gtacctgcag	gcgtaccagc	tttccctata	780
gtgagtcgta	ttagagg					797

<210> 22

<211> 714

<212> DNA

<213> Glycine sp

<220>

<221> misc_feature

<222> (1)...(714)

<223> n = A,T,C or G

<400> 22

aattcggctc	gagacggctg	cgagaagacg	acagaagggg	gttaccgtta	tcatgcaagc	60
tgataatggg	gcctctgaag	ttcttggtcc	gttattataa	aactgagtc	ttcactctct	120
ctcgaaccag	ctcacagaaa	caatgatctc	ctacgccgac	atgctcaagg	gatcacatgg	180
atgtcaacaa	cttcaactat	cctccattgt	cagagatgta	aactacagct	gtggctcgtg	240
tggttatgag	ctgaacttga	actccagcaa	ccgcaacact	tggtctctca	ttgactcaaa	300
gtccataaaag	agaggcatca	tctccttctt	ctccgtggat	gagagcaggt	tcactcagat	360
ccagcaactt	cactggcctt	cttggatgcc	ctttttcaac	tccaagcgcc	aaagaaccaa	420
gcttttttgc	cgcagctgtg	ggaaccacct	tggttatgct	tacactttgc	ctctcaatct	480
caatccccggg	atggcatctc	tgatgattca	gaatctatga	tatcaaaacta	accgctttgt	540
taccttcttt	ctgcgaggaa	ccaagtcaaa	agtttaganga	tatgggcaag	gtttgagact	600
gcatcttcct	ccactcttgg	tggtctaatt	cttgaaaggg	acagaaacat	attcatcagt	660
tcttggttgg	ttggaatgng	aattaatgna	ttctaccttt	tgacattatg	aagg	714

<210> 23

<211> 525

<212> DNA

<213> Glycine sp

<400> 23

cggtctcgag	attactggag	gaaaggacat	aaccctgcag	gaagtgaaca	gggtttctca	60
ggttgtgact	agtttggtg	atccttctgc	taatattata	tttggggctg	tcgttgatga	120
tcgtctacact	ggggagattc	acgtgactat	cattgcaact	ggcttctcac	agtcttttca	180
gaagaagttg	ctaacagatc	caagggctgc	aaagctgctt	gacaagggtg	ctgagggccca	240
agaaagcaag	gtagtccctc	ctccccctcaa	gtcctcaaac	aaggttgaat	ctagaccatc	300
cccgcgaaaag	ctcttttttt	agttgcattg	ttctttttac	ccttttttcat	ttttccaatt	360
attattatta	tattatattg	gccgatcaaa	aaaaaaatta	ttatattata	ttgtaggaca	420
caatgatctt	gatgcttaat	taagtgagat	atcattctct	tgatgttctt	tcccctccaa	480
aaaaaaaaaa	aaagggcggc	cgccgactag	tgagctcgctc	gaccc		525

<210> 24

<211> 1083

<212> DNA

<213> Glycine sp

<400> 24

cggtctgagg	aaccctatta	aaattggaga	agttctgact	cgtggattag	gtacgggagg	60
gaatccactt	ttgggggaac	aagctgcaga	ggaatcaaga	gatgctattg	ctgatgctct	120
taaaggatca	gatttgggtg	ttataacggc	tgggatgggt	gggggaaccg	ggctctgggtgc	180
tgccccagtt	gtagcccaaa	tatcaaaaga	ggcaggttac	ttgactgtag	gtgttggttac	240
ctatcccttc	agttttgaag	gacgtaagag	atccttgcag	gcctttgaag	ccatcgaaaag	300
gctgcagaaa	aatgttgaca	cacttatagt	gattccaaat	gaccgtctgc	ttgacatagc	360
tgatgagcag	atgcctcttc	aggatgcttt	tccgtcttgc	agatgacgtt	ctacggcaag	420
gagtacaggg	aatatcagac	attatamctg	wcctggactt	gtcaatgtgg	atttttgcag	480
atgtaaaagc	tgtgatgaaa	gactctggga	ctgcaatgct	tggagtaggt	gtttcctccg	540
gtaaaaaccg	agcagaagaa	gcagccsaac	aggctacttt	ggctycttta	attggatcct	600
ctatttcagt	cagctactgg	gggtagtgtg	taataattact	ggaggaaagg	acataaccct	660
scaggaagtg	aacagggktt	ctcaggttgt	gactaagttt	ggctgatcct	tctgctaata	720
ttatatttgg	ggctgtcgtt	gatgatcgct	acacggggga	gattcacgtg	actatcattg	780
caactggcct	ctcacagtct	tttcagaaga	agttgctaac	agatccaagg	gctgcaaagc	840
tgcttgacaa	ggtggctgag	ggccaagaaa	gcaaggcagt	ccctcctccc	ctcaagtcct	900
caaacaagggt	tgaatctaga	ccatccccgc	gaaagctctt	tttttagttg	catggttcct	960
tttacccttt	ttcatttttc	caattattat	tattatatta	tattggccga	tcaaaaaaaaa	1020
aaaaaaaggg	cggccgccga	ctagttagct	cgtcgaccgc	ggaattaatt	cgggaccgggt	1080
acc						1083

<210> 25

<211> 1335

<212> DNA

<213> Glycine sp

<220>

<221> misc_feature

<222> (1)...(1335)

<223> n = A,T,C or G

<400> 25

cggtctgagg	cccagaacaa	caaaaattgc	tcctcaacgc	ctaagtcgtc	gtttcggttc	60
ggtgagatgc	tectacgctt	acgtagataa	cgccaaaatt	aaggttgtcg	gcatcggtcgg	120
tggcggcaac	aatgccgtta	atcgcatgat	cggagtggtg	ttgcagggtg	tagacttcta	180
tgcgataaat	accgatgctc	aggcactatt	aaattctgct	gctgagaacc	ctattaaaaat	240
tggagaagtt	ctgactcgtg	gattaggtac	aggtgggaat	ccacttttgg	gggaacaagc	300
tgcgaggaaa	tccagagatg	ctattgctga	tgctcttaaa	ggatcagatt	tggtattttat	360
aacggctggg	atgggtgggg	gaaccgggtc	ttggtgctgc	cccagttgta	gnccaaatat	420
caaaaagaggc	aggntacttt	gactgtaggt	gttggtacct	atcccttcag	ttttgaagga	480
cgtaagagat	gcttgcaggc	ctttgaagcc	atcgaaaggc	tgcaaaaaaa	tgttgcacac	540
ttatagtatt	tccaaatgat	cgtctgcttg	acatancttg	atgaaccaga	tgctatttca	600
aggatgcttt	ycgytytkca	rawkatgtty	tamcgsaarg	sgkacaggga	atatcaagac	660
attwtaacag	gtacctggac	ttgtmaatgt	agattttgct	gatgtaaaam	ctgkgataaa	720
gacttctggg	actgcaatgc	ttggtgtagg	tgtttcatcc	ggtaaaaccg	accagaagaa	780
gcagcagaac	agggtacttt	tggctccttt	aattggatca	tctattcagt	cagctactgg	840
ggtagtgtat	aatattactg	gaggaaagga	cataaccctg	caggaagtga	acagggtttc	900
tcagggtggtg	actagtttgg	ctgatccttc	tgctaataatt	atatttggag	cttggtgttg	960
atgatcgctt	acactgggga	gattcacgtg	actataattg	caactggcct	ctcacagtct	1020
tttcagaaga	agttgctaac	agatccaagg	gctgcaaagc	tgcttgacaa	agtggtctgag	1080
ggccaagaaa	gcaaggcagt	ccctcctccc	cccaagtcct	caatcaagggt	tgaatctaga	1140
ccatccccgc	gaaagctctt	tttgtagttg	catgggtcct	ttaccctttt	cttttttcca	1200
attattatat	tgtaagtcac	tctgtagtac	aatgatcttg	atgcttaatt	tagtgagata	1260
tcattctctt	gatgttaaaa	aaaaaaaaaa	aaaaaaaaaa	aaagggcggc	cgccgactag	1320

tgagctcgtc gaccc

1335

<210> 26

<211> 902

<212> DNA

<213> Glycine sp

<220>

<221> misc_feature

<222> (1)...(902)

<223> n = A,T,C or G

<400> 26

aatcggctc	gagtaccagg	gttgggtgaat	gtagattttg	ctgatgttcg	ggctataatg	60
gccaatgcag	gttcttcact	aatggggata	ggaactgcaa	ctggaaaatc	aagggcaaga	120
gatgctgcat	taaatgccat	ccagtcacct	ttactggata	ttggtatara	gagggctact	180
kgaattgttt	ggaacawaac	tggtgggact	gatctgrcct	tgtttgaggt	aaacacggca	240
gcagagggtta	tttatgacct	cgtggaccct	actgctaatt	taatatattg	agcagtaata	300
gatccatcac	tcagtgggtca	agtggagcata	acattaattg	cttactgrat	tcaaagcgyc	360
aagaggagag	tgaagggagg	cctctgcagg	ccagtcaact	cactcaagca	gacacaacct	420
tcggcaccaa	ttggcgggtc	tcctctttca	ctgatgggtg	tttgtttgag	ataccagaat	480
tcctaagaa	garaggaggt	tcacgctatc	cgagggcgta	atctttttca	tcctaatttc	540
ttttgatccc	ttgcatttct	tcacccttgg	atatacatag	caattgggtc	agttcttarg	600
tcctgtctct	gscctttttc	ggatttwrkc	aaragttgkg	katacagttk	gttcatgaaa	660
gtttattact	tyccactgkc	cagactttatg	ggkctaaacc	gganggtatt	ksarcagga	720
tgcttttctt	ggcatatttg	aattagttta	ttagcttgta	cagagatttc	agtaatgctg	780
agagcttggt	atagttcttt	ggcatgttat	agaaaattca	ttattattaa	aaaaaaaaaa	840
aaagggcgcc	gccgactagt	gagctcgctc	accgggaat	tnattccgga	ccggtacctg	900
ca						902

<210> 27

<211> 856

<212> DNA

<213> Glycine sp

<220>

<221> misc_feature

<222> (1)...(856)

<223> n = A,T,C or G

<400> 27

aattcggctc	gagattgggtg	aaccgtagac	tttgctgatg	ttcgagctat	aatggccaat	60
gcagggtctt	cacttatggg	gataggaact	gcaactggga	aaacaagggc	marggawgct	120
gcattaaatg	ctatccagtc	mccctttact	ggatatttgg	tataraaaagg	gctactggaa	180
ttgtatggaa	cataacyggk	ggaagtgatt	tgaccttgtt	tgaaggtaaa	tggtgcasca	240
raagtatat	atgmccctgt	ggmccccact	gstaatttaa	tatttgggsc	agwaatagat	300
ccatcactcc	agtgggcaag	taagcatamm	wtaatcgcaa	ctggattcaa	gcgtcaagag	360
gaaaagtgaa	gggagaccct	atgcaggcca	gtcaactcac	acaaggagat	nccgttggtg	420
tcaatcggcg	atyttctact	ttcactgatg	gtagcttttg	ttggagatcc	ctggaattct	480
taaagaagaa	ggggcgctca	cgttatccaa	gagtttaata	ctcttttccc	caactcctta	540
atccctcctt	gcatctcttt	mccaascaat	ttttagggat	acaaatctca	tcagtctaag	600
gtattagatc	acgggtttttg	cccccttttt	catttttagg	ttcgcatgtg	gcantamagt	660
tgttcattga	aagcgaagtt	actttccaaa	accgttggtt	tctgarttga	aggcttggtt	720
ggcatgtttt	wataagttta	ttagcttgta	tttttgnca	gagaataata	tatcagtaat	780
ggtcagtgtc	tgttataaan	ccncnaaaaa	aaaaaaaaaa	aaaagggcgg	ccgccgacta	840

gtgagctcgt cgaccc

856

<210> 28
 <211> 1060
 <212> DNA
 <213> Glycine sp

<220>
 <221> misc_feature
 <222> (1)...(1060)
 <223> n = A,T,C or G

```

<400> 28
aattcggctc gaggtcacaa cccctttttc atttgaaggc cgaagaaggc cagttcaagc      60
acaagaagga attgctgcat taagagataa tggtgataca ctgatagtta ttccaaatga      120
caaactgctg actgcagttt ctcaatctac ccctgtaact gaagcattca atctggctga      180
tgatattctt agacaagggtg ttcgtggtat atctgatatt attacgatac caggattggt      240
gaatgtagac tttgcagatg ttcgagctat aatggccaat gcaggttctt cacttatggg      300
gataggaact gcaactggca aaacaagggc aagagacgct gcattaaatg ctatccagtc      360
acctttacta gatattggta tagaaaaggc taccggaatt gtatggaaca taactgggtg      420
aaagwgattt gaccttgttt gaggtaaatg ctgcagcaga agttatata gaccttgtgg      480
accccactgy taatttaata tttggagcag taatagatcc atcactcagt ggtcaagtaa      540
gcatcacatt aattgcaact ggattcaagc cgtcaagagg aaaagtgaag ggagacctat      600
acaggccarc caatttacac aaggagatac ggttggtatc attcttaag aaccsgcgat cttcctctt      660
cactgatggk agctttgttg agayccctga tcttaatccc ttgatttctt taaaaagtaa      720
tccgagagct taatactctt ctccccaatt tcttaatccc ttgatttctt taaaaagtaa      780
tttttaggga tacaatctc atcagctctg gtattagatc ccgttttgcc cttttttttt      840
ttcattttta ggttcgcatt gggcatactg ttgttcaaga aagcaaagta ctttcaaac      900
cgttgtttac tgagtcgagg cttgttgcca ggttttaata agtttattag cttgtatttt      960
ttgtacagag aatatatcag taatggtcag ggcttggtat nnnanncccn nnnannaaan      1020
aaaaaaaaag gcggccgccg actagtgagc tcgtcgaccc      1060

```

<210> 29
 <211> 727
 <212> DNA
 <213> Glycine sp

<220>
 <221> misc_feature
 <222> (1)...(727)
 <223> n = A,T,C or G

```

<400> 29
atctcncaaa atgcatgnen ctgtgtgtgg catatattca aaatgacttg gcccgagggtg      60
gggttttant ttgctcttag aaaatgtgtt gagcctgcac atanaagatt ggagttgttg      120
attctcagtg gattgttcac caagggtatt cctcactagg gaatcagggt gantctcaaa      180
caggaaagcn ccatggcagg ggntgagggg ncggtgtana aaggagtggc catgttccag      240
agtcgggtggc aaatgctgaa tacgcgtatc acaactccat tggaattgat acatctaatt      300
ccactgctca ttaggtgact tcggcctaag ttgacttgta aacatattgt tactaccctt      360
agccttacgc gtagaatttt cccttaaaaa aaaaaatata ttcttatgta acgttacgta      420
catgcaatgc aatcacataa tagagtccta gctagggacc aaacatcatt tcgatgtaga      480
aattgctgta cttaacagtg agtaaactta gtgaagagaa ttattattgc tgctaacgaa      540
ggtgcttata ggaaatggaa atgctagtga atccttaaat tggaggctga caacgaagtt      600
ctttaggggt tttgggatta aagaaaaacga aatgtcataa ttatcatacc cttgggatga      660
ggagacagga ctattactat aaaaaaaaaa aaaaaggggc gccgccgact agtgagctcg      720

```

727

tcgacccc

<210> 30

<211> 1185

<212> DNA

<213> Glycine sp

<220>

<221> misc_feature

<222> (1)...(1185)

<223> n = A,T,C or G

<400> 30

cggtctcgagc	tggaatgggt	gggggaactg	gcacaggtgg	agctccaatt	attgctagta	60
ttgcaaagtc	aatgggtata	ttgacggttg	gtattgtcac	cacccctttc	tcgtttgaag	120
ggagaaagag	atctattcaa	gcccaagaag	gaattacagc	cttaagagat	aatgttgaca	180
cgcttatagt	tattccaaat	gacaagctac	taacggcagt	ttctcaatct	acccctgtaa	240
ctgaagcatt	caatctggct	gatgatattc	ttcgacaggg	tggtcgtggc	atatctgata	300
ttattacaat	accagggttg	gtgaatgtag	atthttgctga	tggtcgggct	ataatggcca	360
atgcaggttc	ttcactaatg	gggataggaa	ctgcaactgg	aaaatcaagg	gcaagagatg	420
ctgcattaaa	tgccatccag	tcaccwttmc	tggtatattg	tatagararg	gctactggaa	480
ttgtttggaa	cawaactggk	gggactgatc	ttgacctgtg	ttgaggtaaa	cacggcarca	540
rraggttatt	tatgacctcg	tggaacctac	tgctaattta	atatttggag	cagtaataga	600
tccatcactc	agtggkcaag	tgagcataac	attaattgct	actggattca	agcgtcaaga	660
ggarartgaa	rggaggcctn	tcagggccag	tcaactcact	caagcagaca	caaccttcgg	720
caccaattgg	cggtcttcct	ctttcactga	tggtggtttg	tttgagatac	cagaattcct	780
aaagaagaga	ggaggttcac	gctatccgag	ggcgtaactc	ttttcactct	aatttctttg	840
atcccttgca	tttcttcacc	cttgatata	catagcattg	gtctagttct	taggtccctg	900
tcttgcctct	tttcggattt	tagtcagagt	tgtgtatata	gtttgttcat	gaaagtttat	960
tacttcccac	tgtccagact	tatgggtcta	accggaggta	ttgcagcatg	gatgcttttc	1020
ttggcatatt	tgaattagtt	tattagcttg	tacagagatt	tcagtaatgc	tgagagcttg	1080
ttatagttct	ttggcatggt	atagaaaatt	cattattatt	attcatcccn	ccaaaaaaa	1140
aaaaaaaaaa	aaagggcggc	cgccgactag	tgagctcgtc	gaccc		1185

<210> 31

<211> 700

<212> DNA

<213> Glycine sp

<220>

<221> misc_feature

<222> (1)...(700)

<223> n = A,T,C or G

<400> 31

aattcggtc	gagattgtca	ccaccccttt	ctcgtttgaa	gggagaaaaga	gatctattca	60
agcccaagaa	ggaattacag	ccttaagaga	taatgttgac	acgcttatag	ttattccaaa	120
tgacaagcta	ctaacggcag	tttctcaatc	taccctgtga	actgaagcat	tcaatctggc	180
tgatgatatt	cttcgacagg	gtgggtccgtg	gcataatctga	tattattaca	ataccagggg	240
tggtgaatgt	agatthttgct	gatgtttcggg	ctataatggc	caatgcagg	tcttcaactaa	300
tggggatagg	aactgcaact	ggaaaatcaa	gggcaagaga	tgctgcatta	aatgccatcc	360
agtcaccttt	actggatatt	ggtatagaga	gggtactgg	aattgthttg	aacataactg	420
gtgggactga	tctgccttgt	ttgaggtaaa	cacngcagca	ganggtattt	atgacctcgn	480
ggccctactg	ctaattaata	tttggagcag	aatagatcca	tcctcatggc	aagtgcata	540
cattnantgc	tctggattca	agcgtcaaga	ngagaagtga	agggangcct	ttgcaggcca	600

gcactcactc agcagacaca accttngnac caattggcgg cttcctcttt cactgatggg 660
 nggttggttg agatncnana attcctaaag aaaaaanagag 700

<210> 32

<211> 1425

<212> DNA

<213> Arabidopsis sp

<400> 32

ccacgcgtcc	ggaggaagta	aacaatggcg	ataattccgt	tagcacagct	taatgagcta	60
acgattttctt	catcttcttc	ttcgtttctt	accaaatega	tatcttctca	ttcgttgcat	120
agtagctgca	tttgcgcaag	ttctagaatc	agtcaattcc	gtggcgggctt	ctctaaacga	180
agaagcgatt	caacaaggct	taagtcgatg	cgattgaggt	gttccttctc	tccgatggaa	240
tctgcgagaa	ttaagggtgat	tggtgtcggt	gggtggtgta	acaatgccgt	taaccggatg	300
atttcaagcg	gtttacagag	tggtgatctc	tatgcgataa	acacggattc	gcaagctctg	360
ttacagtttt	ctgtcgagaa	cccacttcaa	attggagaac	ttttaactcg	tgggcttggc	420
actggtggaa	acccgcttct	tggaagaaca	gctgcagaag	aatcaaaaga	tgcaattgct	480
aatgctctta	aaggatcaga	ccttggtttc	ataactgctg	gtatgggtgg	tggaaacagg	540
tctggtgctg	cacctgtggt	agctcagatt	tcgaaggatg	ctggttattt	gactgttggg	600
gttggttacct	atccggttag	ctttgaagga	cgtaaaagat	ctttgcaggc	actggaagct	660
attgaaaagc	tcctaaaaga	tggtgatacc	cttatcgatg	ttccaaatga	tcgtctgcta	720
gatattgctg	atgaacagac	gccacttcag	gacgcgtttc	ttcttgcaag	tgatgtttta	780
cgccaaggag	tacaagggaat	ctcagatatt	attactatac	ctggactagt	caatgtggat	840
tttgcatgat	tgaaggcagt	catgaaagat	tctggaactg	caatgctcgg	ggtaggtggt	900
tcttccagca	aaaaccgggc	agaagaagca	gctgaacaag	caactttggc	tccattgatc	960
ggatcatcca	tacaatcagc	tactggtgtc	gtctacaaca	tcactggtgg	aaaagacata	1020
actttgcagg	aagtgaaccg	agtatcacag	gtcgtgacaa	gtttggcaga	cccacgggac	1080
aacatcatat	ttggagctgt	tggtgatgat	cgctacaccg	gagagattca	tgtaacgata	1140
atcgccacag	gcttctctca	gtcattccag	aagacacttc	tgactgatcc	aagagcagct	1200
aaactccttg	acaaaatggg	atcatcaggt	caacaagaga	acaaagggaat	gtctctgcct	1260
caccagaagc	agtctccatc	aactatctct	accaaactcg	cttctccccg	tagacttttc	1320
ttctagtttt	cttttttttc	tttctggttt	caagcatcaa	aaatgtaacy	atcttcaggc	1380
tcaaatatca	attacatttg	attttctctc	aaaaaaaaaa	aaaaa		1425

<210> 33

<211> 1611

<212> DNA

<213> Arabidopsis sp

<400> 33

tggtgttgcc	gctcagaaat	ctgaatcttc	tccaatcaga	aactctccac	ggcattacca	60
aagccaagct	caagatcctt	tcttgaacct	tcacccggaa	atatctatgc	ttagaggtga	120
agggactagt	acaatatgca	atccaagaaa	ggaaacgtct	tctggacctg	ttgtcgagga	180
ttttgaagag	ccatctgctc	cgagtaacta	caatgaggcg	aggattaagg	ttattggtgt	240
gggagggtgt	ggatcaaatg	ctgtgaatcg	tatgatagag	agtgaatgt	cagggtgtgga	300
gttctggatt	gtcaacactg	atatccaggc	tatgagaatg	tctcctgttt	tgccgtgataa	360
taggttacaa	attggttaag	agttgactag	gggttttaggt	gctggaggaa	atccagaaat	420
cggtatgaat	gctgctagag	agagcaaaga	agttattgaa	gaagctcttt	atggctcaga	480
tatggtcttt	gtcacagctg	gaatgggcgg	tggaactggc	actggtgcag	cccctgtaat	540
tgcaggaatt	gccaaggcga	tggttatatt	gacagttggt	attgccacaa	cgcttttctc	600
gtttgagggg	cgaagaagaa	ctgttcaggc	tcaagaaggg	cttgcatctc	tcagagacaa	660
tggtgacact	ctcatcgta	ttccaaatga	caagttgctt	acagctgtct	ctcagctctac	720
tccggttaaca	gaagcattta	atctagctga	tgatatactc	cgtcaggggg	ttcgtgggat	780
atctgatata	attacgattc	ctgggttggt	gaatgtggat	tttgctgatg	tgagagctat	840
aatggcaaat	gcgggggtctt	cattgatggg	aataggaact	gcgacaggaa	agagtcgggc	900

```

aagagatgct gcgctaaatg caatccaatc ccctttgtta gatattggga ttgagagagc 960
cactgggaatt gtttggaaca ttactggcgg aagtgacttg acattgtttg aggtaaatgc 1020
tgctgcggaa gtaatatatg atcttgtcga tccaactgcc aatcttatat tcgtgctggt 1080
gtagatccag ccctcagcgg tcaagtaagc ataaccctga tagctacggg tttcaaacga 1140
caagaagagg gagaaggacg aacagttcag atgggtacaag cagatgctgc gtcagttgga 1200
gctacaagaa gaccctcttc ttcctttaga gaaagcgggt cagtggagat cccagagttc 1260
ttgaagaaga aaggcagctc tcgttatccc cgagtctaaa gcccaatcta atcactacct 1320
tgcacactgc agcaataaca aacgtgtgtg tactggtagt ctggtagtgc cttctgggat 1380
acagcaagat gtgttgatgt atgatcaaga atctgtgtgg gtgtgtatat gttctgtcac 1440
tgctctgtgt cgtgttcttg aataggttgt tttagaaatc ggagtttctc tctatgtcac 1500
ttccaaaaca aaaaaggaga agaagaatca cacttctcga accataaaca tacttataag 1560
attatgagag ttttagcaga aattattgtc aaaaaaaaaa aaaaaaaaaa a 1611

```

<210> 34
 <211> 299
 <212> DNA
 <213> Arabidopsis sp

<220>
 <221> misc_feature
 <222> (1)...(299)
 <223> n = A,T,C or G

```

<400> 34
agtaattgaa aaatgacaca tacctctttt nctacaatca aagtttataa ctaagaagaa 60
caaatgcaat agtaataata aagattttgag gacatactgt gacaaagacc atatctgagc 120
cataaagagc ttctncaata actgctttgc tctctctagc agcattcata ccgatttcng 180
gatttctctc agcacctaaa cccctagtca actccttacc aatttgtaac ctattatcag 240
gcaaaacagg agacattctc atagcctgga tatcagtgtt gacaatccag aactccaca 299

```

<210> 35
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic Oligonucleotide

```

<400> 35
cttgccgcaa aacatcatcc gcgag 25

```

<210> 36
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic Oligonucleotide

```

<400> 36
caccaccccc gagcatcgca gttccgga 28

```

<210> 37
 <211> 41
 <212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 37

ttggtgtcgg tgggtggtggt aacaatgccg ttaaccggat g

41

<210> 38

<211> 41

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 38

ggtaacaatg ccgtaaccg gatgatttca agcggtttac a

41

<210> 39

<211> 41

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 39

gctgctcttg gatcagtcag aagtgtcttc tggaatgact g

41

<210> 40

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 40

tttgtcaagg agtttagctg ctcttggatc agtcagaag

39

<210> 41

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 41

cagcaccaga tcctgtacct ccacccat

28

<210> 42

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 42

ctgaatgggt atgtgacaac accaacagtc

30

<210> 43

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 43

gctgatgatg tattacgcca aggtgtcc

28

<210> 44

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 44

ctggaactgc tatgcttgga gttggggt

28

<210> 45

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 45

tcgaaactga atggataggt gacaacacca acag

34

<210> 46

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 46

tggatcgaca aggtcgtaga taatttcggc cgcagc

36

<210> 47

<211> 56

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 47

cgcgatttaa atggcgcgcc ctgcaggcgg ccgcctgcag ggcgcgccat ttaa

56

<210> 48

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 48

tcgaggatcc gcggccgcaa gcttcctgca gg

32

<210> 49

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 49

tcgacctgca ggaagcttgc ggccgcggat cc

32

<210> 50

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 50

tcgacctgca ggaagcttgc ggccgcggat cc

32

<210> 51

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 51

tcgaggatcc gcggccgcaa gcttcctgca gg

32

<210> 52

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 52

tcgaggatcc gcggccgcaa gcttcctgca ggagct

36

<210> 53

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 53

cctgcaggaa gcttgcggcc gcggatcc

28

<210> 54

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 54

tcgacctgca ggaagcttgc ggccgcggat ccagct

36

<210> 55

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 55

ggatccgcgg ccgcaagctt cctgcagg

28

<210> 56

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> c-myc Tag

<400> 56

Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu
1 5 10

<210> 57

<211> 37

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 57
acgtgcggcc gcatggcgat aattccgtta gcacagc

37

<210> 58
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 58
actgataagc ttttgctcga agaaaagtct acggggga

37

<210> 59
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 59
acgtgcggcc gcatggcgat aattccgtta gcacagc

37

<210> 60
<211> 44
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 60
cgtcctgcag gctacaagtc ttcctcactg ataagctttt gctc

44

<210> 61
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 61
acgtcctgca ggatggcgat aattccgtta gcacagc

37

<210> 62
<211> 38

<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 62
acgtgcggcc gcctagaaga aaagtctacg gggagaag 38

<210> 63
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 63
acgtaagctt tccttctctc cgatggagtc tg 32

<210> 64
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 64
acgtctgcag cttttcaatg gcttcaagtc cct 33

<210> 65
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 65
acgtctgcag aagaacgtgg ataccctcat cgtg 34

<210> 66
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 66
acgtgcggcc gcctagaaga acaatctacg gggagaag 38

<210> 67
<211> 244

<212> DNA

<213> Artificial Sequence

<220>

<223> Prnr/NEP/Gene10leader/14 amino acid from GFP fusion

<400> 67

gaattcggta	ccccgcgtcgt	tcaatgagaa	tgataagag	gctcgtggga	ttgacgtgag	60
ggggcaggga	tggttatatt	tctgggagcg	aactccgggc	gaatactgaa	gcgcttggat	120
acaagttatc	cttgggaagga	aagacaattc	cggatcctct	agaaataatt	ttgtttaact	180
ttaagaagga	gatataccca	tggttaaagg	agaagaactt	ttcactggag	ttgtcccaag	240
catg						244

<210> 68

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 68

gtcctgcagg	atggccacca	tctcaaacc	ag	32
------------	------------	-----------	----	----

<210> 69

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Oligonucleotide

<400> 69

tagcgccgc	ctatataaag	gagctaaaag	aacagcc	37
-----------	------------	------------	---------	----

<210> 70

<211> 669

<212> DNA

<213> Arabidopsis sp

<400> 70

atggcgatta	gtccgttggc	acagcttaac	gagctaccag	tctcttctc	gtttsttgcg	60
acatcccact	cgctgcacag	taccagaatc	agtggcgggc	ttctcaaac	aaaggtttaa	120
gcaaacacgg	ttgagatgct	ccttctctcc	gatggagtct	gcgaggatta	aggtggttgg	180
tgctcgcggt	ggtggtaaca	atgccgtcaa	tcgcatgatt	tccagcggct	tacagagtgt	240
tgatttctat	gcgataaaca	cggactctca	agctctcttg	cagtcttctg	cgcagaaccc	300
tcttcaaatt	ggagagctcc	taactcgtgg	ccttgggact	ggtgggaacc	cgcttctagg	360
agaacaagct	gctgaggaat	ctaaagacgc	gattgctaata	gctcttaaac	gatctgacct	420
tgytttcatt	actgctggta	tggttggtgg	cactggctcc	ggtgctgctc	ctgttggtgc	480
tcagatctcc	aaagacgctg	gttatttgac	cgttggtggt	gttacctatc	ccttcagctt	540
cgaaggtcgt	aaaagatctt	tgcaggcact	tgaagccatt	gaaaagctgc	agaakaacgt	600
ggataccctc	atcgtgatac	caaatgatcg	tctcctagat	attgctgatg	aacagacgcc	660
tcttcaaga						669

<210> 71
 <211> 646
 <212> DNA
 <213> Nicotiana sp

```

<400> 71
ggccctctag atgcatgctc gagcggccgc cagtgtgatg gatatctgca gaattcgccc      60
ttaagcagtg gtaacaacgc agagtacgcg ggggttaaacc aaacagacag agagagcaga      120
aacagcaatg gccaccatct caaaccacgc agagatagca gcttcttctc cttcctttgc      180
tttttaccac tcttccttta ttcctaaaca atgctgcttc accaaagctc gccggaaaag      240
cttatgtaaa cctcaacggt tcagcatttc aagttcattt actccttttg attctgctaa      300
gattaaggtt atcggcgctc gtggcggtgg taacaatgcc gttaaccgta tgattggcag      360
tggtcttacag ggtgttgact tctatgctat aaacacggat gctcaagcac tgctgcagtc      420
tgctgctgaa aatccacttc aaattggaga gcttctgact cgtgggcttg gtactgggtg      480
caatcctctt ttaggggaac aggcagcaga ggagtcgaag gaagccattg caaattctct      540
aaaaggttca gatatggtgt tcataacagc aggaatgggt ggaggtacag gatctggtgc      600
tgaagggcga attccagcac actggcggcc gttactagtg gatccg                      646
  
```

<210> 72
 <211> 325
 <212> PRT
 <213> Glycine sp

```

<400> 72
Gly Ser Arg Pro Arg Thr Thr Lys Ile Ala Pro Gln Arg Leu Ser Arg
 1      5      10      15
Arg Phe Gly Ser Val Arg Cys Ser Tyr Ala Tyr Val Asp Asn Ala Lys
 20      25      30
Ile Lys Val Val Gly Ile Gly Gly Gly Asn Asn Ala Val Asn Arg
 35      40      45
Met Ile Gly Ser Gly Leu Gln Gly Val Asp Phe Tyr Ala Ile Asn Thr
 50      55      60
Asp Ala Gln Ala Leu Leu Asn Ser Ala Ala Glu Asn Pro Ile Lys Ile
 65      70      75      80
Gly Glu Val Leu Thr Arg Gly Leu Gly Thr Gly Gly Asn Pro Leu Leu
 85      90      95
Gly Glu Gln Ala Ala Glu Glu Ser Arg Asp Ala Ile Ala Asp Ala Leu
 100     105     110
Lys Gly Ser Asp Leu Val Phe Ile Thr Ala Gly Met Gly Gly Gly Thr
 115     120     125
Gly Ser Gly Ala Ala Pro Val Val Ala Gln Ile Ser Lys Glu Ala Gly
 130     135     140
Tyr Leu Thr Val Gly Val Val Thr Tyr Pro Phe Ser Phe Glu Gly Arg
 145     150     155     160
Lys Arg Ser Leu Gln Ala Phe Glu Ala Ile Glu Arg Leu Gln Lys Asn
 165     170     175
Val Asp Thr Leu Ile Val Ile Pro Asn Asp Arg Leu Leu Asp Ile Ala
 180     185     190
Asp Glu Gln Met Pro Leu Gln Asp Ala Phe Pro Phe Ala Asp Asp Val
 195     200     205
Leu Arg Gln Gly Val Gln Gly Ile Ser Asp Ile Ile Thr Val Pro Gly
 210     215     220
Leu Val Asn Val Asp Phe Ala Asp Val Lys Ala Val Met Lys Asp Ser
 225     230     235     240
Gly Thr Ala Met Leu Gly Val Gly Val Ser Ser Gly Lys Asn Arg Ala
  
```

```

                245                250                255
Glu Glu Ala Ala Glu Gln Ala Thr Leu Ala Pro Leu Ile Gly Ser Ser
                260                265                270
Ile Gln Ser Ala Thr Gly Val Val Tyr Asn Ile Thr Gly Gly Lys Asp
                275                280                285
Ile Thr Leu Gln Glu Val Asn Arg Val Ser Gln Val Val Thr Ser Leu
                290                295                300
Ala Asp Pro Ser Ala Asn Ile Ile Phe Gly Ala Val Val Asp Asp Arg
305                310                315                320
Tyr Thr Gly Glu Ile
                325

```

<210> 73
 <211> 357
 <212> PRT
 <213> Zea mays

<220>
 <221> VARIANT
 <222> (1)...(357)
 <223> Xaa = Any Amino Acid

```

<400> 73
Asp Leu Val Phe Ile Thr Ala Gly Met Gly Gly Gly Thr Gly Ser Gly
1          5          10
Ala Ala Pro Val Val Ala Gln Ile Ser Lys Glu Ala Gly Tyr Leu Thr
20          25          30
Val Gly Val Val Thr Tyr Pro Phe Ser Phe Glu Gly Arg Lys Arg Ser
35          40          45
Val Gln Ala Leu Glu Ala Leu Glu Lys Leu Glu Lys Ser Val Asp Thr
50          55          60
Leu Ile Val Ile Pro Asn Asp Lys Leu Leu Asp Val Ala Asp Glu Asn
65          70          75          80
Met Pro Leu Gln Asp Ala Phe Leu Leu Ala Asp Asp Val Leu Arg Gln
85          90          95
Gly Val Gln Gly Ile Ser Asp Ile Ile Thr Ile Pro Gly Leu Val Asn
100         105         110
Val Asp Phe Ala Asp Val Lys Ala Val Met Lys Asn Ser Gly Thr Ala
115         120         125
Met Leu Gly Val Gly Val Ser Ser Ser Lys Asn Arg Ala Gln Glu Ala
130         135         140
Ala Glu Gln Ala Thr Leu Ala Pro Leu Ile Gly Ser Ser Ile Glu Ala
145         150         155         160
Ala Thr Gly Val Val Tyr Asn Ile Thr Gly Gly Lys Asp Ile Thr Leu
165         170         175
Gln Glu Val Asn Lys Val Ser Gln Ile Val Thr Ser Leu Ala Asp Pro
180         185         190
Ser Ala Asn Ile Ile Phe Gly Ala Val Val Asp Asp Arg Tyr Thr Gly
195         200         205
Glu Ile His Val Thr Ile Ile Ala Thr Gly Phe Pro Gln Ser Phe Gln
210         215         220
Lys Ser Leu Leu Ala Asp Pro Lys Gly Ala Arg Ile Val Glu Ser Lys
225         230         235         240
Glu Lys Ala Ala Thr Leu Ala His Lys Ala Ala Ala Ala Val Gln
245         250         255

```

Pro	Val	Pro	Ala	Ser	Ala	Trp	Ser	Arg	Arg	Leu	Phe	Ser	Xaa	Glu	Ala
			260					265					270		
His	Leu	Val	Asn	Arg	Asp	Ser	Xaa	Cys	Ile	Arg	Phe	Ala	Phe	Ser	Val
		275					280					285			
Leu	Arg	Ala	Val	Pro	Lys	Val	Ile	Phe	Gly	Tyr	Leu	Glu	Ile	Tyr	Ser
	290					295					300				
Leu	Gly	Xaa	Cys	Ser	Val	Val	Val	Glu	Xaa	Val	Ser	Val	Tyr	Val	Ser
	305				310					315					320
Leu	Leu	Cys	Phe	Met	Phe	Leu	Arg	Ile	Xaa	Arg	Xaa	Gly	Xaa	Glu	Lys
				325					330					335	
Cys	Ser	Ala	Thr	Gln	His	Xaa	Thr	Val	Xaa	Lys	Ile	Phe	Asp	Cys	Phe
			340					345					350		
Ile	Ala	Ala	Thr	Cys											
			355												

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/28103

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/82 C12N15/29 C07K14/415 C12N5/10 A01H1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	<p>WO 98 00436 A (UNIV NEVADA) 8 January 1998 (1998-01-08)</p> <p>abstract; claims 1,8 page 2, line 30 -page 3, line 24 page 6, line 3 - line 16 page 6, line 33 -page 7, line 4 page 7, line 21 -page 8, line 31 page 10, line 26 - line 36</p> <p>--- -/--</p>	<p>16, 17, 24-26, 30 1-11, 18-23</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

4 April 2000

Date of mailing of the international search report

17/04/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ceder, O

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/28103

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	STREPP ET AL: "Plant nuclear gene knockout reveals a role in plastid division for the homolog of the bacterial cell division protein FtsZ, an ancestral tubulin" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 95, 1 April 1998 (1998-04-01), pages 4368-4373, XP002083808 ISSN: 0027-8424 cited in the application	16, 24-26, 30
A	abstract page 4368, left-hand column	6-12
X	SVAB ET AL: "High-frequency plastid transformation in tobacco by selection for a chimeric aadA gene" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 90, February 1993 (1993-02), pages 913-917, XP002106110 ISSN: 0027-8424 cited in the application	27
A	abstract	1
A	YU ET AL.: "Inhibition of assembly of bacterial cell division protein FtsZ by the hydrophobic dye 5,5'-bis-(8-anilino-1-naphthalenesulfonate)" THE JOURNAL OF BIOCHEMICAL CHEMISTRY, vol. 273, no. 17, 24 April 1998 (1998-04-24), pages 10216-10222, XP002134449 cited in the application abstract page 10221, right-hand column, line 5 - line 7 page 10221, right-hand column, line 25 -page 10222, left-hand column, line 3	6, 12-14
A	WO 93 10253 A (ESCAPGENETICS INC) 27 May 1993 (1993-05-27) abstract; claim 1 page 5, line 14 - line 17	1, 2

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/28103

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9800436 A	08-01-1998	AU 3644497 A	21-01-1998
		CA 2259209 A	08-01-1998
		EP 0912596 A	06-05-1999
		US 5981836 A	09-11-1999
WO 9310253 A	27-05-1993	US 5344775 A	06-09-1994
		AU 3139593 A	15-06-1993

